

MELISSA

The Melittologist's Newsletter



Ronald J. McGinley, Bryan N. Danforth, Maureen J. Mello

Department of Entomology • Smithsonian Institution, NHB-105 • Washington, DC 20560

NUMBER — 5

Spring, 1992

COLLECTING NEWS


Report on the first PCAM expedition to northern Mexico

By

Jerome Rozen, Jr.

Department of Entomology,
American Museum of Natural History,
Central Park at 79th St., New York, NY 10024

The first PCAM expedition to Mexico, supported by the U.S. National Science Foundation, explored the northern, arid regions of Mexico as well as the Sierra Madre Occidental, separating the Sonoran and Chihuahuan Deserts.



The trip began on August 12 in Hermosillo, Sonora, (see map) and included Ricardo Ayala (Estación de Biología, Universidad Autónoma de México [UNAM], Chimalpa), Alberto Bürquez (Centro de Ecología, UNAM, Hermosillo), Luis Godínez (Centro de Ecología, UNAM), Terry Griswold (Bee Biology and Systematics Lab., USDA), Bob Minckley (University of Kansas), Jack Neff (Central Texas Melittological Institute), Adrián Quijada (Centro de

Ecología, UNAM), Dr. and Mrs. Wallace LaBerge (Illinois Natural History Survey) and Nancy Pember (American Museum of Natural History). The trip was organized and lead by Alberto Bürquez and myself. Jack Neff collected and pressed all plant specimens, which were subsequently identified at the University of Texas, Austin, thanks to the efforts of Dr. Berle Simpson and her staff.

For the first four days, as the group participants gathered in Hermosillo, collecting trips were taken to the west of Hermosillo, to the Sea of Cortez, and in the mountains to the east of Hermosillo (see map). Unfortunately, flowering in the lowlands of Sonora had been delayed because the late summer rainy season had started only shortly before our arrival. The mountains to the east-northeast of Hermosillo (toward Moctezuma), however, provided excellent collecting. On a single day trip along the Rio Sonora 1,200 specimens were collected. Other collecting highlights around Hermosillo included finding *Calliopsis anomoptera* and *C. limbus*, two *Euphorbia* bees not previously collected so far south.

On August 16 the group departed eastward from Hermosillo to Yécora, Sonora, where specimens of *Thygater cockerelli* were collected. This is the first record of this genus from Sonora and the first record west of the Continental Divide. This species was fairly common further east, in Cuahtémoc. The collection of *Bombus ephippatus* at Yécora also extends the range of this common Mexican bumblebee into Sonora. Specimens of *Colletes platycnema*, the largest member of the genus in the New World, were also collected here. Perhaps the most interesting collections at Yécora were specimens of the predominantly tropical meliponine genus *Partamona*.

Further east, in Chihuahua, we found new species of *Conanthalicus* and *Eufriesia* at General Trias. The magnificent, all metallic-blue *Eufriesia* was collected visiting yellow-flowered *Solanum*, and extends the range of this tropical genus far to the north. A new species of *Calliopsis* (*Perissander*) was also collected

near Villa Ahumada, Chihuahua. In general, collecting in the Chihuahuan desert was excellent due to record rainfalls for the season.

At Ciudad Jiménez the LaBerges and Nancy Pember returned to the United States and the remaining participants continued on to Reserva Biosfera, Mapimi, a remote site on the border of Durango and Chihuahua. This was undoubtedly the most successful collecting locality of the entire trip — more than 4000 specimens were collected over several days. At least one, and possibly three, new species of *Melanomada* were collected here, as well as *Calliopsis (Micronomadopsis) meliloti* and *Anthocopa daleae*, both previously un-recorded species from Mexico.

Alberto Burquez and Adrián Quijada returned to Hermosillo from Mapimi and the group continued south, reaching the southern-most extent of the trip, in the vicinity of Torreón and Zapata, on August 24. *Calliopsis (Micronomadopsis) callosa*, which had previously been known only from western Texas and southern Arizona/New Mexico, was collected at Zapata.

Returning northward, the group travelled to Ciudad Chihuahua, with a brief sortie eastward to Ojinaga, on the Rio Grande. A new species of *Stelis*, two new species of *Holcopasites*, and *Deltoptila* sp. were collected on this leg of the trip. We found excellent collecting at the Samalayuca sand dunes south of Ciudad Juarez. Due to abundant flowering of many plants we collected several thousand specimens of genera typical of the southwestern U.S., including a large matinal, *Oenothera*-visiting *Megachile (Megachiloidea)* sp. Males and females of *Perdita (Macropteropsis) magniceps* were collected here; a species oligoleptic on *Sphaeralcea* which had previously been known only from the male holotype.

After Samalayuca the group headed westward with the intent of taking the road from Nuevo Casas Grandes to Cuahtémoc, and on to Yécora. However, heavy rains had washed out the road to Cuahtémoc and we were forced to follow the highway south of the U.S./Mexico border, passing through Agua Prieta, and then west-southwest to Hermosillo. After spending the night of September 1 in Hermosillo, the group, rejoined by Alberto Bürquez, went southward to Alamos, near the southern border of Sonora. *Eulaema polychroma* was collected here, extending the northern-most record of this large genus of orchid bees. The group returned to Hermosillo on September 4 and most participants returned to their respective institutions on September 6.

In summary, the trip was a success. Roughly 17,300 bee specimens were netted during the 26 day, 6,177 km expedition. 148 herbarium specimens were also collected.

* * *

Report on the second NSF-Funded PCAM expedition

By

Terry Griswold
USDA Bee Biology and Systematics Laboratory,
Utah State University, UMC 53,
Logan, Utah 84322-5310

The second PCAM expedition to Mexico, supported by NSF grant BFR-9024723 with assistance from the Instituto de Biología, UNAM, was conducted 28 October-13 November 1991 to explore the late fall fauna of parts of the Altiplano and coastal Veracruz with emphasis on the xeric areas of the Tehuacan Desert, Puebla and Metztitlan Valley, Hidalgo — areas thought by some to be desert refugia during the Pleistocene. Participants for the entire expedition were Ricardo Ayala, Felipe Noguera, and Terry Griswold, with Alicia Rodríguez and Catalina Everaert joining for parts of the trip.

The expedition began in Mexico City, moved south and east to Tlayacapan, Taxco, and Ixtapan de la Sal, then southeast to Acatlán and Huajuapan de León, northeast through the Tehuacan Desert to Xalap, down to Cardel, then north along the coast to Poza Rica, inland to Pachuca, north through the Metztitlan Valley to Otango, and returned to Mexico City. The final two days were spent collecting in areas just south of the city and in natural area on the campus of UNAM.

The area sampled included parts of the states of Mexico, Morelos, Guerrero, Oaxaca, Puebla, Veracruz, and Hidalgo, and the Distrito Federal. Parts of the route traveled are probably the most completely sampled of any region in Mexico. But collections during the summer months account for almost all of the sampling effort; very few bees have been collected during fall months. The results of this expedition prove that this is not due to the absence of a fall bee fauna. Over 4600 specimens were obtained during fifty collector-days. Bees were abundant in a wide variety of habitats including pine and oak forests, cloud forest, subtropical and tropical forests, and deserts. Except in a few areas, such as the coastal dunes of Veracruz, flowers were abundant. Some of the best collecting was on Asteraceae, Fabaceae, And *Salvia*.

Seventy-seven genera of bees were collected. Abundant groups included *Andrena*, Panurginae, Megachilidae, Eucerini, and *Bombus*. Surprisingly, Colletidae and Halictini were very poorly represented. Notable finds include the following: 1) Numerous specimens of the rare anaphoroid *Deltoptila*; 2) The first records of *Dufourea* south of Zacatecas, *D. cyanella* from Morelos and Guerrero and an undescribed species from Puebla; 3) A new species of *Hesperapis* from the Tehuacan Desert; 4) A nesting site of the rarely collected *Mesoxaea tachytiformis*; 5) Seven

species of *Lasioglossum*, two of them apparently new; 6) Several new species of *Protandrena*; 6) Good series of *Trachusa* (*Ulanthidium*) and *T.* (*Heteranthidium*); 7) A new species of *Megachile* (*Chelostomoides*) from Ixtapan de la Sal, Mexico; 8) A new species of *Osmia* (*Diceratosmia*); 9) An undescribed species of *Hoplitis* (*Eremosmia*) from the desert area of Metztitlan Valley; 10) An undescribed osmiine of questionable generic assignment near *Hoplitis* or *Asmeadiella*, also from the Metztitlan Valley; 11) The previously undescribed female of *Nomada* (*Pachynomada*) *dreisbachorum*; 12) An undescribed nomadine belonging to the *Melanomada* complex; and 13) A new species of *Syntrichalonia*. Of interest was the relatively large number of specimens with bent hairs on the clypeus and/or frons of the female. These included *Caupolicana* (*Zikanapis*) *clypeata*, three species of *Lasioglossum*, *Osmia* (*Diceratosmia*) n.sp., *Anthidium*, *Trachusa* (*Utanthidium*), *Megachile* (*Chelostomoides*) n.sp., *Exomalopsis binotata*, *Pectinapis*, *Deltoptila*, and *Habropoda* aff. *salviarum*. Almost all these species were collected on *Salvia*.

A further expedition at this season or even later should be profitable. Flowers at many of the sites did not appear to be past their peak. Most of the specimens collected were in good condition, with the possibility of additional species later in the year. Areas which were poorly sampled due to weather or simply the time constraints of the trip included the road between Ixtapan de la Sal and Chalma, the road between Tehuacan and Orizaba, the eastern slopes of Perote and Citlaltepetl, the road from Poza Rica to Tulancingo, and the Metztitlan Valley. Areas where flowering was mostly past which should be sampled in September include the road between Izucar de Matamoros and Huajuapan, the Tehuacan Desert, xeric areas of Puebla around Perote, the coastal dunes of Veracruz, and the road between Tulancingo and Pachuca.

March PCAM Trip to Coahuila

By

Robert Brooks and Byron Alexander
Snow Entomological Museum, University of Kansas,
Lawrence, KS 66045

The next collecting trip in the faunal survey of Mexican bees (PCAM) will take place from 13 March to 3 April. Robert Brooks and Byron Alexander (University of Kansas) are in charge of arrangements in the U.S., whereas Ricardo Ayala (Estacion de Biologia, Universidad Autonoma de Mexico, Chamela) will be handling arrangements in Mexico. Jack Neff (Central Texas Melittological Institute) will be accompanying us as a combined bee specialist and botanist, along with Luis Godinez (UNAM) and Doug Yanega (University of Kansas). We plan to take a single vehicle, the VW van recently purchased with grant funds.

Our goal primary goal is to survey the inland "Coahuila Chaparral," a habitat with a restricted, patchy distribution in northern Mexico. Chaparral is one of the ecological zones that the grant specifically targets and so most of the time will be spent visiting the various Sierras of this region, that range up to 2200 m in elevation. With these objectives in mind, we are planning to spend most of the three-week trip in the state of Coahuila, primarily north of Cuatro Cienegas.

Participants will gather in Torreon, and then travel northward to the Reserva Biosfera in Mapimi, a rich collecting site visited on previous PCAM expeditions. From there we will go further north to Sierra La Madera, just west of Cuatro Cienegas, and on to Ocampo, where we will leave the pavement and will try and reach Sierra El Fuste, Sierra San Ambrosio, Sierra El Palomino, Sierra Mojada, Sierra El Colorado and Sierra La Encantada.



RESEARCH NEWS

A Revised Checklist of Polish Colletidae and Halictidae

By

Tadeusz Pawlikowski
Copernicus University,
Torun, Poland

During the last 15 years of my investigations on the Polish Apoidea, I have made many local lists of bees collected in Poland over the last century (see *Melissa* 2: 12-13).

The status of the generic categories are as in Michener C.D. (1944, *Bull. Amer. Mus. Nat. Hist.* 82: 151-326; 1964, *Amer. Zool.* 4: 227-239). However the species are in alphabetical order within each genus. The species records are divided into two date classes: pre-1950 (+), and 1950 to the present (*).

A list containing 34 species of Colletidae and 101 species of Halictidae from Poland is presented below.

COLLETIDAE

Hylaeus Fabricius, 1793

= *Prosopis* Fabricius, 1804

- * *angustatus* (Schenck, 1859)
- * *annularis* (Kirby, 1802)
- = *cervicornis* (Costa, 1858)
- = *dilatata* (Kirby, 1802)
- * *annulatus* (Linnaeus, 1758)
- = *borealis* Nylander, 1852
- + *bisinuatus* Förster, 1871
- = *leptocephala* (Morawitz, 1871)
- * *brevicornis* Nylander, 1852
- * *cypearis* (Schenck, 1853)
- * *communis* Nylander, 1852
- * *confusus* Nylander, 1852

- = *gibba confusus* Nylander
 sq. Warncke, 1972
 + *comutus* Curtis, 1831
 * *diformis* (Eversmann, 1852)
 + *gibbus* Saunders, 1850
 = *genalis* Thompson, 1872
 = *mixta* (Schenck, 1859)
 * *gracilicornis* (Morawitz, 1867)
 * *hyalinatus* Smith, 1842
 * *nigritus* (Fabricius, 1798)
 * *pectoralis* Förster, 1871
 = *kriechbaumeri* Förster, 1871
 = *palustris* Perkins auct.
 * *pictipes* Nylander, 1852
 * *punctatus* (Brullé, 1832)
 * *punctulatissimus* Smith, 1842
 + *rinki* (Gorski, 1852)
 * *signatus* (Panzer, 1798)
 = *pratensis* Geoffroy auct.
 * *sinuatus* (Schenck, 1853)
 = *minutus* (Fabricius, 1793)?
 * *styriacus* Förster, 1871
 + *variegatus* (Fabricius, 1798)
- Colletes* Latreille, 1802
 * *caspicus balticus* Alfken, 1912
 * *cunicularius* (Linnaeus, 1761)
 * *daviesanus* Smith, 1846
 = *inexpectatus* Noskiewicz, 1936
 * *floralis* Eversmann, 1852
 = *montanus* Morawitz, 1876
 * *fodiens* Fourcroy, 1785
 * *impunctatus* Nylander, 1852
 = *alpinus* Morawitz, 1872
 * *marginatus* Smith, 1846
 * *nasutus* Smith, 1853
 * *punctatus* Mocsary, 1877
 * *similis* Schenck, 1853
 = *pistigma* Thomson, 1872
 * *succinctus* (Linnaeus, 1758)
- HALICTIDAE**
- Halictus* Robertson, 1918
 * *confusus perkinsi* Blüthgen, 1926
 * *eurygnathus* Blüthgen, 1931
 * *leucaheneus arenosus* Ebmer, 1976
 = *fasciatus* auct. nec Nylander, 1848
 * *maculatus* Smith, 1848
 * *quadricinctus* (Fabricius, 1776)
 * *rubicundus* (Christ, 1791)
 * *semitectus* Morawitz, 1873
 * *sexcinctus* (Fabricius, 1793)
 * *simplex* Blüthgen, 1923
 * *subauratus* (Rossi, 1792)
 * *tetrazonius* (Klug, 1817)
 * *tumulorum* (Linnaeus, 1758)
 = *fasciatus* Nylander, 1848
- LasioGLOSSUM* Curtis, 1833
 * *aeratum* (Kirby, 1802)
 = *viridiaeneum* (Blüthgen, 1918)
 * *albipes* (Fabricius, 1781)
 * *bavaricum* (Blüthgen, 1930)
 * *breviceps* (Schenck, 1868)
 + *breviventre* (Schenck, 1853)
 * *calceatum* (Scopoli, 1763)
 = *cylindricum* (Fabricius, 1763)
 * *convexusculum* (Schenck, 1853)
- * *costulatum* (Kriechbaumer, 1873)
 + *cupromicans* (Pérez, 1903)
 * *euboense* (Strand, 1909)
 * *fratellum* (Pérez, 1903)
 = *niger* (Viereck, 1903)
 * *fulvicorne* (Kirby, 1802)
 * *glabriusculum* (Morawitz, 1872)
 * *intermedium* (Schenck, 1868)
 * *interruptum* (Panzer, 1798)
 * *laeve* (Kirby, 1802)
 * *laevigatum* (Kirby, 1802)
 * *laterale* (Brullé, 1832)
 * *laticeps* (Schenck, 1868)
 = *mendax* (Alfken, 1912)
 * *lativentre* (Schenck, 1853)
 * *leucopum* (Kirby, 1802)
 * *leucozonium* (Schrank, 1781)
 * *limbellum* (Morawitz, 1876)
 * *lineare* (Schenck, 1868)
 * *lucidulum* (Schenck, 1861)
 * *major* (Nylander, 1852)
 * *malachurum* (Kirby, 1802)
 = *longulum* (Smith, 1848)
 * *minutissimum* (Kirby, 1802)
 * *minutulum* (Schenck, 1853)
 * *morio* (Fabricius, 1793)
 * *nigripes* (Lepeletier, 1841)
 * *nitidisculum* (Kirby, 1802)
 * *nitidulum aenidorsum* (Alfken, 1921)
 * *nitidulum nitidulum* (Fabricius, 1804)
 = *continentale* (Blüthgen, 1944)
 = *smeathmanellus* auct. nec Kirby, 1802
 * *parvulum* (Schenck, 1853)
 = *minutum* (Kirby, 1802) nec (Fabricius, 1798)
 * *pauillum* (Schenck, 1853)
 * *politum* (Schenck, 1853)
 = *nanulum* (Schenck, 1853)
 * *prasimum haemorrhoidale* (Schenck, 1853)
 * *punctatissimum* (Schenck, 1853)
 + *puncticolle* (Morawitz, 1872)
 + *pygmaeum* (Schenck, 1853)
 * *quadrinotatum* (Schenck, 1861)
 * *quadrinotatum* (Kirby, 1802)
 * *quadrivirgatum* (Schenck, 1853)
 * *rufitarse* (Zetterstedt, 1838)
 * *semilucens* (Alfken, 1914)
 * *setulellum* (Strand, 1909)
 * *setulosum* (Strand, 1909)
 + *sexmaculatum* (Schenck, 1853)
 * *sexnotatum* (Nylander, 1852)
 * *sexnotatum* (Kirby, 1802)
 = *nitidum* Panzer, 1798
 * *sexstrigatum* (Schenck, 1868)
 * *subfasciatum* (Imhoff, 1832)
 = *rufocinctum* (Nylander, 1852)
 * *tarsatum* (Schenck, 1868)
 * *tricinctum* (Schenck, 1874)
 * *vilosulum* (Kirby, 1802)
 * *xanthopum* (Kirby, 1802)
 * *zonulum* (Smith, 1848)
- Sphecodes* Latreille, 1804
 * *albilabris* (Fabricius, 1793)
 = *fuscipennis* (Germar, 1819)
 * *crassus* Thomson, 1870
 = *variegatus* Hagens, 1882
 * *cristatus* Hagens, 1882

- * *croaticus* Meyer, 1922
- * *ephippius* (Linneaus, 1767)
- = *divisus* (Kirby, 1802)
- = *rufescens* (Geoffroy in Fourcroy, 1785)
- = *similis* Wesmael, 1865
- * *fasciatus* Hagens, 1882
- = *affinis* Hagens, 1882
- * *ferruginatus* Hagens, 1882
- * *gibbus* (Linnaeus, 1758)
- * *hyalinatus* Hagens, 1882
- * *longulus* Hagens, 1882
- * *marginatus* Hagens, 1882
- * *miniatus* Hagens, 1882
- * *monilicornis* (Kirby, 1802)
- = *subquadratus* Smith, 1845
- * *niger* Hagens, 1874
- * *pellucidus* Smith, 1845
- = *pilifrons* Thomson, 1870
- * *puriticeps* Thomson, 1870
- * *reticulatus* Thomson, 1870
- + *rubicundus* Hagens, 1875
- * *rufiventris* (Panzer, 1798)
- = *subovalis* Schenck, 1853
- + *scabricollis* Wesmael, 1865
- + *spinulosus* Hagens, 1875
- Nomioides* Schenck, 1866
 - * *minutissimus* (Rossi, 1790)
- Dufourea* Lepetier, 1841
 - = *Halictoides* Nylander, 1848
 - * *dentiventris* (Nylander, 1848)
 - + *halictula* (Nylander, 1848)
 - * *inermis* (Nylander, 1848)
 - * *vulgaris* (Nylander, 1859)
- Rophites* Spinola, 1808
 - = *Rhophitoides* Schenck, 1859
 - * *canus* Eversmann, 1852
 - * *hartmani* Friese, 1902
 - * *quinquespinosus* Spinola, 1808
 - + *trispinosus* Pérez, 1903
- Systropha* Illiger, 1806
 - * *curvicornis* (Scopoli, 1770)
 - * *planidens* Giraud, 1861

Sex Determination in Early Instar Bee Larvae

We wish to bring to the attention of Melissa readers a recent paper describing very simple staining methods for determining the sex of bee larvae at any stage of development, from first instar to pre-pupa. M.J. Duchateau and P. Van Leeuwen (1990, Insectes Sociaux, 37:232-235) describe their use of the method to determine the sex of bumble bee larvae but speculate the method may work on other bees (and presumably wasps) as well.

The method relies on first fixing the larvae in Carnoy's fixative and then using Congo Red to stain the imaginal disks of the genitalia, which differ strikingly in morphology between males and females. The method works very well with *Perdita* last instar larvae, even when the staining step is eliminated. We thank Hayo H.W. Velthuis and George Eickwort for pointing this article out to us.

Phenology of two territorial solitary bees, *Anthidium manicatum* and *A. florentinum* (Hymenoptera: Megachilidae)

By
Peter Witz

Five years' data on phenology of an *Anthidium manicatum* population in Southern Germany and comparative observations on *A. manicatum* and *A. florentinum* from Southern France are analyzed. Males and females had the same flight season, adult sex ratio was strongly female biased and males were larger than females in both species. This is the opposite pattern to most other solitary bees, where females are generally larger than males, sex ratio is male-biased, males emerge before females and males disappear long before females. We argue that two features of *Anthidium* female behaviour, namely prolonged sexual receptivity and use of resources easily defendable by males, explain male adaptations in behaviour, phenology, and body size and, hence population sex ratio. (Full article to be published in Journal of Zoology.)

Research at Tel-Aviv University

Dan Eisikowitch wanted to share with us some of the work going on at the Tel-Aviv University in Israel.

"Our labs are collaborating on a joint project on the intimate relations between *Calotropis procera*, a species of Asclepiadaceae, and the bees on which it depends for pollination, *Xylocopa pubescens* and *X. sulcatipes*.

"This research is supported by the Israeli Academy of Science and its main target is to elucidate the way in which the plant attracts the bees, information about what the bees leave on the flowers and how other bees can evaluate it.

"We shall appreciate comment from anyone interested in such interrelationships."

MEETING ANNOUNCEMENTS

Fifth International Conference on Apiculture in Tropical Climates

September 7 - 12 1992.

Organized by The International Bee Research and Association and The Government of Trinidad and Tobago. The purpose of this meeting is to focus attention on the bees and beekeeping of the tropics and subtropics. For more information please send your name and address to: Nicola Bradbear, Fifth International Conference on Apiculture in Tropical Climates, International Bee Research Association, 18 North Road, Cardiff, CF1 3DY, UK.

International Workshop on Non-*Apis* Bees

August 10-13, 1992

USDA-ARS Bee Biology & Systematics Laboratory

Utah State University

Logan, Utah 84322-5310

Second Announcement and Call for Papers

Monday - August 10

The Biology and Management of the Alfalfa Leafcutting Bee

Moderators: W.P. Stephen, Oregon State University, Corvallis

K.W. Richards, Agriculture Canada, Lethbridge

Tuesday - August 11

Bumblebees as Crop Pollinators

Moderator: R.M. Fisher, Acadia University, Wolfville

Non-Social Bees as Crop Pollinators

Moderator: S.W.T. Batra, USDA ARS, Beltsville

Wednesday - August 12

Foraging and Nesting Behavior of Non-*Apis* Bees

Moderator: P.G. Willmer, University of St. Andrews, Fife

Bee-Plant Interactions

Moderator: A. Shmida, Hebrew University, Jerusalem

Thursday - August 13

Interactions Between Bees and Their Parasites

Moderator: To be announced

Biosystematics, Biogeography and Evolution of Bees

Moderator: C.D. Michener, University of Kansas, Lawrence

Contact: Vincent J. Tepedino (801) 750-2559

Philip F. Torchio (801) 750-2520

MARGINALIA

Chris Starr sent us this item from *Folk Tales of Meghalaya and Arunachal* (1974, Prades, New Delhi: Sterling, 112 pp.) by B.K. Borgohain.

How Bees Came Into the World

The god named Limir-Chabbo looked like a bison. At the time when the sun and moon shed light on the earth in the beginning, Limir-Chabbo roamed about in the mountains. While he was thus roaming about his dropping turned into bees and those bees drank up his urine and so they did not leave the place. Then another god named Dalling picked up the bees and put them on a tree. But they slipped down from the tree. Then the god fixed the hive on the tree by tying it with his loin cloth. The bees built their nest sitting up on the hive and produced honey. The honey is sweet. Because it is god's urine.

CORRECTIONS

In the Recent Literature list included with Melissa 4, two papers, Gonzalez, J.M., 1989 and Gonzalez, J.M., 1990 were given single authorship. They were both co-authored and should have read Gonzalez, J.M. and Gainai, M.A.

The second edition of *Bumblebees* by Oliver Prys-Jones and Sarah Corbet, has been published by the Richmond Publishing Co. Anyone interested in ordering this book should write to the following address: Richmond Publishing Co. Limited, P.O. Box 963, Slough SL2 3RS, England. (ISBN: 0.8356 2574 [paperback]; 0.83546 2582 [hardcover])

Books of interest:

Kin Recognition. Peter G. Hepper, ed. Cambridge University Press, New York, 1991. See the review in Science 255:217.

Recent Apoid Literature: 1987-1991

The following list of references has been compiled primarily from the National Agricultural Library's computer-based bibliographic retrieval system, as well as from reprints or lists of papers kindly sent to us by subscribers. Papers dealing specifically with the commercial aspects of *Apis mellifera* or *Megachile rotundata* have not been included, but papers of general biological interest have been included. Key words included most bee genera, familial and subfamilial names as well as pollination biology and bee/plant interactions. We admit in advance that these citations have not been checked for accuracy.

At present the roughly 500 references shown below are included in a single file created by v. 1.4 of the bibliographic software package Pro-Cite (Address: P.O. Box 4250, Ann Arbor, Michigan, 48106; 313-996-1580). The entire bibliography contains over 1300 references in two files. We would like to make this and expanded lists available to other Pro-Cite users, or users of other, compatible, bibliographic programs. We hope in the future to send out the complete bibliography on disk to readers who agree to add references or check those already input. Additionally, it is planned to supplement this list with references gathered from Biological Abstracts or Zoological Record, which should extend the list back to the late 1970's.

As a preliminary step towards making this reference list available to *Melissa* readers on disk, we would be willing to send copies of the full bibliography in the form of a WordPerfect 5.0 text file to anyone who sends us a request and a blank 3.5 or 5.25 inch diskette.

Please send us comments on the utility of the reference list and how we might make it more generally available to researchers. And please continue to send reprints and reference lists to Ronald McGinley for inclusion in upcoming lists.

Bibliography

- Abrol, D. P. (1988). Effect of climatic factors on pollination activity of alfalfa-pollinating subtropical bees *Megachile nana* Bingh and *Megachile flavipes* Spinola (Hymenoptera: Megachilidae). *Acta Oecologica*, 9(4), 371-377.
- Abrol, D. P. (1990). Pollination activity of alfalfa-pollinating subtropical bees *Megachile nana* and *Megachile flavipes* (Hymenoptera: Megachilidae). *Trop. Ecol.*, 31(1), 106-115.
- Abrol, D. P., & Kapil, R. P. (1988). Foraging range of *Andrena ilerda* and *Andrena leaena* (Hymenoptera: Andrenidae). *Research and Development Reporter*, 5(1/2), 65-67.
- Abrol, D. P., & Kapil, R. P. (1989). Wing beat frequency of alfalfa pollinating bees *Megachile nana* Bingh and *Megachile flavipes* Spinola (Hymenoptera: Megachilidae). *J. Anim. Morphol. Physiol.*, 36(2), 191-194.
- Accorti, M. (1991). Bees and agricultural productivity. *Ethol. Ecol. Evol.*, (Spec. issue 1), 162.
- Albuquerque, C. M. R.D., Hadj-Idirs, A. E. Q., & Hertz, E. (1989). Evolution and perisynaptic organs in the ventral nerve cord of *Melipona scutellaris* Latreille 1811 (Hymenoptera: Apidae: Meliponinae). *Rev. Bras. Entomol.*, 33(3-4), 421-428.
- Albuquerque, P. M. C., & Rego, M. M. C. (1990). Phenology of bees visiting murici trees *Byrsonima crassifolia* Malpighiaceae. *Bol. Mus. Para Emilio Goeldi Ser. Zool.*, 5(2), 163-178.
- Alcock, J. (1990). Body size and territorial behavior in the bee *Protoxaea gloriosa* (Fox) (Hymenoptera: Oxaeidae). *Pan-Pac. Entomol.*, 66(2), 157-161.
- Alcock, J., & Johnson, M. D. (1990). Female mate choice in the carpenter bee *Xylocopa varipuncta* (Hymenoptera: Anthophoridae). *J. Zool. (Lond.)*, 221(1), 195-204.
- Alexander, B. (1990). A cladistic analysis of the Nomadine bees (Hymenoptera: Apoidea). *Syst. Entomol.*, 15(2), 121-152.
- Alexander, B. (1991). *Nomada* phylogeny reconsidered (Hymenoptera: Anthophoridae). *J. Nat. Hist.*, 25(2), 315-330.
- Alexander, B. (1991). Phylogenetic analysis of the genus *Apis* (Hymenoptera: Apidae). *Ann. Entomol. Soc. Am.*, 84(2), 137-149.
- Alexander, B. (1990). A preliminary phylogenetic analysis of sphecid wasps and bees. *Sphecos*, (20), 7-16.
- Aliev, K. A. (1989). Zoogeographic bee complexes (Hymenoptera: Apoidea) in the lesser caucasus within Azerbaijan SSR USSR. *Izv. Akad. Nauk. Az. SSR Ser. Biol. Nauk.*, (3), 54-61.
- Almeida, M. C. D., & Laroca, S. (1988). *Trigona spinipes* (Apidae: Meliponinae) taxonomy bionomy and trophic relationships in restricted areas. *Acta Biol. Parana*, 17(1-4), 67-108.
- Alzuet, A. D. B.D., & Abrahamovich, A. H. (1990). Types of association between Acari and Hymenoptera II. Description of the stages of the life cycle of *Sennertia augustii* new species (Acarina: Chaetodactylidae) associated to *Xylocopa augusti* Lepeletier 1841 (Hymenoptera: Anthophoridae). *Rev. Bras. Entomol.*, 34(3), 627-636.

- Archer, M. E. (1990). The aculeate solitary wasps and bees (Hymenoptera: Aculeata) of Leicestershire. *Trans. Leic. Lit. Philos. Soc.*, 84, 9-25.
- Archer, M. E. (1989). Notable wasps and bees (Hym., Aculeata) taken in the Breckland and Suffolk Sandlings. *Entomol. Mon. Mag.*, 125(Sept-Dec), 232.
- Archer, M. E. (1991). Notable wasps and bees (Hymenoptera: Aculeata) in the Isle of Wight new forest and Hampshire England UK during 1988. *Entomol. Mon. Mag.*, 127, 34.
- Archer, M. E. (1991). Notable wasps and bees (Hymenoptera: Aculeata) taken in Gloucestershire Somerset Devon and Dorset England UK during 1985. *Entomol. Mon. Mag.*, 127, 76.
- Archer, M. E. (1990). Seasonal flight activity of bumblebees (Hymenoptera: Apinae) as monitored by malaise trap catches. *Br. J. Entomol. Nat. Hist.*, 3(4), 173-176.
- Archer, M. E. (1990). The solitary aculeate wasps and bees (Hymenoptera: Aculeata) of an English suburban garden. *Entomologist's Gazette*, 41(3), 129-142.
- Armbruster, W. S. (1988). Multilevel comparative analysis of morphology, function and evolution of *Dalechampia* blossoms. *Ecology*, 69(6), 1746-1761.
- Armbruster, W. S., & McCormick, K. D. (1990). Diel foraging patterns of male euglossine bees: ecological causes and evolutionary response by plants. *Biotropica*, 22(2), 160-171.
- Armbruster, W. S. (1990). Estimating and testing the shapes of adaptive surfaces: the morphology and pollination of *Dalechampia* blossoms. *Am. Nat.*, 135(1), 14-31.
- Armstrong, D. P., & Paton, D. C. (1990). Methods for measuring amounts of energy available from *Banksia* inflorescences. *Aust. J. Ecol.*, 15(3), 291-298.
- Arshad, M., & Parveen, M. (1987). Studies on the poison apparatus of *Megachile lanata* Fabr. *Proc. Pak. Congr. Zool.*, 7, 177-180.
- Ayala, R. (1988). Abejas silvestres (Hymenoptera: Apoidea) de Chamela, Jalisco, Mexico. *Folia Entomol. Mex.*, (77), 395-493.
- Ayasse, M. (1990). Visual and olfactory signals involved in nest-finding of *Lasioglossum malachurum* (Hymenoptera: Halictidae). *Apidologie*, 21(4), 349-351.
- Ayasse, M., Leys, R., Pamilo, P., & Tengo, I. (1990). Kinship in communally nesting *Andrena* (Hymenoptera: Andrenidae) bees as indicated by composition of Dufour's gland secretions. *Biochem. Syst. Ecol.*, 18 (6), 453-460.
- Banaszak, J. (1990). Bees (Apoidea) of hornbeam-oak and thermophilous oak forests of the Mazovian lowland. *Zeszyty Nauk. Wydziały Szkoly Pedagogicznej w Bydgoszczy*, 8, 24-36.
- Batra, S. W. T. (1990). Bionomics of a vernal solitary bee *Andrena (Scrapteropsis) alleghaniensis* Viereck in the Adirondacks of New York (Hymenoptera: Andrenidae). *J. Kansas Entomol. Soc.*, 63(2), 260-266.
- Batra, S. W. T. (1990). Bionomics of *Evylaeus comagenensis* (Knerer and Atwood) (Halictidae), a facultatively polygynous, univoltine, boreal halictine bee. *Proc. Entomol. Soc. Wash.*, 92(4), 725-731.
- Bego, L. R., Maeta, Y., Tezuka, T., & Ishida, K. (1989). Floral preference and flower constancy of a Brazilian stingless bee *Nannotrigona testaceicornis* kept in a greenhouse (Hymenoptera: Apidae). *Bull. Fac. Agric. Shimane Univ.*, (23), 46-54.
- Bilinski, M. (1988). The optimum time for transferring *Megachile rotundata* nests for wintering. *Pszczelnicze Zeszyty Naukowe*, 32, 89-98.
- Bischoff de Alzuet, A., & Abrahamovich, A. H. (1991). Description of the life cycle stages of *Sennertia splendidulae* Alzuet and Abrahamovich 1989 (Acarina: Chaetodactylidae) associated with *Xylocopa splendidula* Lepeletier 1841 (Hymenoptera: Anthophoridae). *Stud. Neotrop. Fauna Environ.*, 26(1), 1-10.
- Bogdanov, S., Imdorf, A., Kilchermann, V., & Gerig, L. (1990). Ruckstände von Folbex (VA) in Wachs, Futter und Honig. *Schweizerische Bienen-Zeitung*, 113(5), 250-254.
- Bonelli, B., & Campadelli, G. (1989-90). Note biologiche su *Chalicodoma parietina* Geoffr. (Hymenoptera: Megachilidae). *Boll. Ist. Ent. << G. Grandi>> Univ. Bologna*, 44, 1-9.
- Boomsma, J. J. (1991). Adaptive colony sex ratios in primitively eusocial bees. *Trends Ecol. Evol.*, 6(3), 92-95.
- Borg-Karlsson, A. K. (1990). Chemical and ethological studies of pollination in the genus *Ophrys* (Orchidaceae). *Phytochemistry*, 29(5), 1359-1387.

- Bull, J. J., & Rice, W. R. (1991). Distinguishing mechanisms for the evolution of cooperation. *J. Theor. Biol.*, 149(1), 63-74.
- Bullock, S. H., Ayala, R., Rodriguez-Zamora, D., Quiroz-Garcia, D. L., & Luz Arreguin-Sanchez, M. D. de la. (1991). Nest provision and pollen foraging in three mexican species of solitary bees (Hymenoptera: Apoidea). *Pan-Pac. Entomol.*, 67(3), 171-176.
- Bullock, S. H., del Rio, C. M., & Ayala, R. (1989). Bee visitation rates to trees of *Prockia crucis* differing in flower number. *Oecologia*, 78, 389-393.
- Camargo, J. M. F. (1988). Meliponinae (Hymenoptera, Apidae) da colecao do Istituto di Entomologia Agraria, Portici, Italy. *Revista Brasileira de Entomologia*, 32(3/4), 351-374.
- Camargo, J. M. F., & Moure, J. S. (1990). Two new species of *Lestrimelitta* Friese (Meliponinae: Apidae: Hymenoptera) from the Amazon region Brazil. *Bol. Mus. Para Emilio Goeldi Ser. Zool.*, 5(2), 195-212.
- Camargo, J. M. F., & Posey, D. A. (1991). Knowledge of the kayapo on stingless social bees (Meliponidae, Apidae, Hymenoptera) additional notes. *Bol. Mus. Para Emilio Goeldi Ser Zool.*, 6(1), 17-42.
- Camargo, J. M. F., & Wittmann, D. (1989). Nest architecture and distribution of the primitive stingless bee, *Moureella caerulea* (Hymenoptera, Apidae, Meliponinae): Evidence for the origin of *Plebeia* (S. Lat.) on the Gondwana continent. *Stud. Neotrop. Fauna Environ.*, 24(4), 213-229.
- Cameron, S. A. (1990). Mitochondrial DNA evolution in bees: single or multiple origins of highly social behavior. In: *Proceedings of the Fourth International Congress of Systematic and Evolutionary Biology*; College Park, Maryland, USA, July 1-7, 1990.
- Cameron, S. A., & Robinson, G. E. (1990). Juvenile hormone does not affect division of labor in bumble bee colonies (Hymenoptera: Apidae). *Ann. Entomol. Soc. Am.*, 83(3), 626-631.
- Campbell, D. R. (1991). Comparing pollen dispersal and gene flow in a natural population. *Evol.*, 45(8), 1965-1969.
- Campos, L. A. de Oliveira, & Costa, M. A. (1990). Determinacao do sexo em abelhas 28. Determinacao das castas em *Schwarziana quadripuncta* (Hymenoptera, Apidae). *Rev. Bras. Biol.*, 49(4), 999-1001.
- Campos, L. A. de Oliveira, & de Melo, G. A. R. (1990). Physogastric-queen mating in *Melipona quadrifasciata* Lep. (Hymenoptera, Apidae). *Rev. Bras. Genet.*, 13(3), 491-500.
- Cane, J. H., & Buchmann, S. L. (1989). Novel pollen-harvesting behavior by the bee *Protandrena mexicanorum* (Hymenoptera: Andrenidae). *J. Insect Behav.*, 2(3), 431-436.
- Cane, J. H., & Payne, J. A. (1990). Native bee pollinates rabbiteye blueberry. *Highlights Agric. Res. Ala. Agric. Exp. Stn.*, 37(4), 4.
- Cartar, R. V. (1991). Colony energy requirements affect response to predation risk in foraging bumble bees. *Ethology*, 87(1/2), 90-96.
- Cartar, R. V. (1991). A test of risk-sensitive foraging in wild bumble bees. *Ecol. Publ. Ecol. Soc. Am.*, 72(5), 888-895.
- Cartar, R. V., & Dill, L. M. (1990). Colony energy requirements affect the foraging currency of bumble bees. *Behav. Ecol. Sociobiol.*, 27 (5), 377-383.
- Cartar, R. V., & Dill, L. M. (1991). Costs of energy shortfall for bumble bee colonies: predation, social parasitism, and brood development. *Can. Entomol.*, 123(2), 283-293.
- Castillo, R. F. del, & Gonzalez-Espinoza, M. (1988). An evolutionary interpretation of sexual polymorphism in *Opuntia robusta*. *Agrociencia, Mexico*, (71), 185-196.
- Castro, L. (1989). Sobre los Bombinae (Hymenoptera: Apidae) de las Sierras de Cazorla, segura y alcaraz. *Anales de Biol.*, 15, 95-100.
- Celary, W. (1989). Review of the parasitic bees of the family Megachilidae (Hymenoptera, Apoidea) in Poland. *Pol. Pismo. Entomol.*, 59(2), 335-355.
- Celary, W. (1989). Review of the cuckoo bees of the genus *Psithyrus* Lep. (Hymenoptera, Apidae) in Poland. *Ann. Upper Silesian Mus. Bytom. Nat. Hist.*, (12), 67-74.
- Celli, G., Porrini, C., & Raboni, F. (1989). Monitoraggio con api della presenza dei ditiocarbammati nell'ambiente (1983-1986). *Boll. 1st Entomol. Univ. Studi. Bologna*, 43, 195-205.
- Chapman, T. W., Wilsie, A. P., Kevan, P. G., & Willis, D. S. (1990). Fiberglass resin for determining nest architecture of ground nesting bees. *J. Kansas Entomol. Soc.*, 63(4), 641-643.
- Chen, G. P., & et al. (1989). Bionomics of *Megachile (Megachile) subtriquilla* Yasumatsu and its control. *Kunchong Zhishi*, 26(4), 219-220.

- Chen, H. M., & Chang, Y. Z. (1987). The structural differentiation of mouthparts of leaf-cutting bees and their biological significance. *Acta Agric. Univ. Pekinensis*, 13(1), 69-76.
- Chiappa, T. E., Rojas, G.-L. M., & Toro, G. H. (1990). Key for the bee genera of Chile (Hymenoptera: Apoidea). *Rev. Chil. Entomol.*, 18, 67-82.
- Christofferson, L. J., & Brander, P. E. (1990). Pollination of woody landscape plants in greenhouses by leafcutter bees. *Tidsskrift for Planteavl.*, 94(2), 191-194.
- Clifford, P. T. P., & Scott, D. (1989). Inflorescence, bumble bee, and climate interactions in seed crops of a tetraploid red clover (*Trifolium pratense* L.). *J. Applied Seed Production*, 7, 38-45.
- Cmak, J., & Szczypciak, J. (1987). The bumble-bees, *Bombus* of the Holy Cross Mountains National Park. *Chronmy. Przr. Ojczysta.*, 43(5-6), 60-64.
- Coelho, J. R. (1991). Heat transfer and body temperatures in honey bee (Hymenoptera: Apidae) drones and workers. *Environ. Entomol.*, 20(6), 1627-1635.
- Colln, K. (1990). Die Hummeln und Schmarotzerhummeln von Gondersdorf (Kr. Daun). Beitrage zur Insektenfauna der Eifeldorfer 2. *Dendrocopos*, (17), 109-117.
- Cresswell, J. E. (1990). How and why do nectar-foraging bumblebees initiate movements between inflorescences of wild bergamot *Monarda fistulosa* (Lamiaceae)? *Oecologia (Berl)*, 82(4), 450-460.
- Crisanto, M., & Tonesi, R. (1989). Records of *Bombus* in Lombardy (Italy). *Apicultura*, (5), 47-61.
- Cropper, S. C., & Calder, D. M. (1990). The floral biology of *Theelmitra epipactoides* (Orchidaceae), and the implications of pollination by deceit on the survival of this rare orchid. *Plant Syst. Evol.*, 170(1-2), 11-27.
- Cross, E. A., & Bohart, G. E. (1990). Notes on the life history of *Sarcassania boharti* (Acar: Acaridae) and its relationship to the alkali bee, *Nomia melanderi* (Hymenoptera: Halictidae). *J. Kansas Entomol. Soc.*, 63(4), 603-610.
- Cruz-Landim, C. D., & Giometti, S. E. E. (1991). Histolysis of the thoracic larvae muscles during the metamorphosis and myogenesis of the flight muscles in *Melipona quadrifasciata anthidioides* (Hymenoptera: Apidae: Meliponinae). *Rev. Bras. Biol.*, 50 (4), 983-1002.
- Cure, J. R., & Wittmann, D. (1990). *Callonychium petuniae* a new panurgine bee species (Apoidea, Andrenidae) oligolectie on *Petunia* (Solanaceae). *Stud. Neotrop. Fauna Environ.*, 25(3), 153-156.
- d'Albore, G. R., & Roscioni, T. (1990). Pollinators of sainfoin *Onobrychis viciifolia* Scop. in mountain environment. *Apic. Mod.*, 81(5), 195-201.
- da Cunha, M. A. S., Cruz-Landim, C. D., & Campos, L. A. D. O. (1990). Occurrence and morphology of the epidermal glands in some species of bees (Hymenoptera: Apoidea). *Naturalia*, 15, 209-217.
- da Silveira, F. A. (1991). Influence of pollen grain volume on the estimation of the relative importance of its source to bees. *Apidologie*, 22(5), 495-502.
- Danforth, B. N. (1991). Female foraging and intranest behavior of a communal bee, *Perdita portalis* (Hymenoptera: Andrenidae). *Ann. Entomol. Soc. Am.*, 84(5), 537-548.
- Danforth, B. N. (1991). The morphology and behavior of dimorphic males in *Perdita portalis* (Hymenoptera: Andrenidae). *Behav. Ecol. Sociobiol.*, 29, 235-247.
- Darakjian, P. (1989). Biologia geral de *Melipona mandacaia*. *Apicultura & Polinizacao*, (31), 24-25.
- Das, M. K., Bandyopadhyay, A. K., & Pawar, A. D. (1990). New host and locality records of *Halictophagus bipunctatus* Yang (Strepsiptera: Halictophagidae) on *Nephrotettix virescens* (Distant) from West Bengal, India. *J. Advanced Zool.*, 10(1), 68-69.
- de Albuquerque, C. M. R., Hadj-Idris, A. E. Q., & Hertz, E. (1989). Evolution and perisynaptic organs in the ventral nerve cord of *Melipona scutellaris* (Latreille, 1811) (Hymenoptera, Apidae, Meliponinae). *Rev. Bras. Entomol.*, 33(3-4), 421-428.
- de Almeida, M. C., & Laroca, S. (1988). *Trigona spinipes* (Apidae, Meliponinae): Taxonomia, Bionomia e relacoes troficas em areas restritas. *Acta Biologica do Parana*, 17(1/4), 67-108.
- Dequech, S. I. B., & Becker, M. (1990). Polinizacao de alfalfa (*Medicago sativa* L.). 1. Especies de abelhas visitantes da cultura. *An. Soc. Entomol. Bras.*, 19(2), 423-435.
- Dimitrov, P. (1990). Effect of the density of wild bees (Hymenoptera, Apoidea) on percentage pollination of lucerne flowers. *Rasteniev DNI Nauki*, 27(6), 23-27.

- Dimitrov, P. (1989). Study on the population dynamics of wild bees (Hymenoptera, Apoidea) in lucerne grown for seed production. Rasteniev DNI Nauki, 26(2), 62-67.
- Donath, H. (1989). Bumble bees as pollinators of lucerne crops. Biol. Stud. Luckau., 18, 57-60.
- Doroshina, L. P. (1990). Protection from chalcids and optimization of nesting conditions for bees (Hymenoptera: Apidoidea: Megachilidae). Entomol. Rev., 69(6), 68-72.
- Dozhdikov, A. A. (1990). Data on the ecology of the nesting site of bumblebees. Ekologiya, (2), 82-84.
- Duchateau, M. J. (1989). Agnostic behaviours in colonies of the bumblebee *Bombus terrestris*. J. Ethol., 7(2), 141-151.
- Duchateau, M. J. (1991). Regulation of colony development in bumblebees. Acta Horticulture 288, 6th Pollination Symposium, p.139-143.
- Duchateau, M. J., & van Leeuwen, P. (1990). Early sex determination in larvae of *Bombus terrestris*. Insectes Soc., 37(3), 232-235.
- Duchateau, M. J., & Velthius, H. H. W. (1988). Development and reproductive strategies in *Bombus terrestris* colonies. Behaviour, 107 (3/4), 186-207.
- Dudley, R., & Ellington, C. (1990). Mechanics of forward flight in bumblebees. 1. Kinematics and morphology. J. Exp. Biol., 148, 19-52.
- Dukas, R., & Dafni, A. (1990). Buzz-pollination in three nectariferous Boraginaceae and possible evolution of buzz-pollinated flowers. Plant Syst. Evol., 169(1-2), 65-68.
- Dylewska, M. (1989). Apoidea of the Ojcow National Park. Part 1. Colletidae, Halictidae, Andrenidae, Melittidae, Megachilidae, Anthophoridae. Acta Biol. Cracov. Ser. Zool., 30, 19-72.
- Eardley, C. D. (1989). The afrotropical species of *Eucera* Friese *Tetralonia* Spinola and *Tetraloniella* Ashmead (Hymenoptera: Anthophoridae). S. Afr. Dep. Agric. Water Supply Entomol. Mem. (Pretoria), (75), 1-62.
- Eardley, C. D. (1991). The genus *Epeolus* Latreille from subsaharan Africa (Hymenoptera: Anthophoridae). J. Nat. Hist., 25(3), 711-732.
- Eardley, C. D. (1991). The Melectini in subsaharan Africa (Hymenoptera: Anthophoridae). S. Afr. Dep. Agric. Dev. Entomol. Mem., (82), 1-48.
- Eardley, C. D., & Brooks, R. W. (1989). The genus *Andrena* Latreille in southern Africa (Hymenoptera: Anthophoridae). S. Afr. Dep. Agric. Water Supply Entomol. Mem., (76), 1-55.
- Eardley, C. D., & Brooks, R. W. (1989). The genus *Anthophora* Latreille in southern Africa (Hymenoptera: Anthophoridae). S. Afr. Dep. Agric. Water Supply Entomol. Mem. (Pretoria), (76), 1-55.
- Eardley, C. D., & Schwarz, M. (1991). The afrotropical species of *Nomada scopoli* (Hymenoptera: Anthophoridae). Phytophylactica, 23(1), 17-28.
- Efremova, Z. A. (1987). The current population of bumblebees of the Volga Region. Dokl. Mosk. O-Vo. Ispyt. Prir. Zool. Bot., 35-38.
- Eijnde, J. v. den. (1990). Method for continuous rearing of *Bombus terrestris* and the production of bumblebees colonies for pollination purposes. Apidologie, 21(4), 330-332.
- Ellington, C. P., Machin, K. E., & Casey, T. M. (1990). Oxygen consumption of bumblebees in forward flight. Nature (Lond), 347(6292), 472-473.
- Ember, A. W., & Sakagami, S. F. (1990). The first record of *Lasioglossum algirum pseudannulipes* new record Bluthgen in Japan with notes on the *Lasioglossum leucopus* group (Hymenoptera: Halictidae). Jpn. J. Entomol., 58(4), 835-838.
- Ember, A. W., & Sakagami, S. F. (1990). *Lasioglossum (Evylaeus) algirum pseudannulipes* (Bluthgen) Ermals in Japan Gefunden, mit Notizen über die L. (E.) *leucopus*- Gruppe (Hymenoptera, Halictidae). Jpn. J. Ent., 58(4), 835-838.
- Engels, W. (1990). Social Insects: An Evolutionary Approach to Castes and Reproduction. Berlin: Springer-Verlag, 264pp.
- Engels, W., Engels, E., Luebke, G., Schroeder, W., & Francke, W. (1990). Volatile cephalic secretions of drones queens and workers in relation to reproduction in the stingless bee *Scaptotrigona postica* (Hymenoptera: Apidae: Trigonini). Entomol. Gen., 15(2), 91-101.
- Engels, W., & Imperatriz-Fonseca, V. L. (1990). Caste development, reproductive strategies, and control of fertility in honey bees and stingless bees. In: W. Engels, Social Insects: An Evolutionary Approach to Castes and Reproduction. (pp. 167-230). Berlin: Springer-Verlag.

- Epperson, B. K., & Clegg, M. T. (1987). Frequency-dependent variation for outcrossing rate among flower-color morphs of *Ipomoea purpurea*. *Evolution*, 41 (6), 1302-1311.
- Evans, H., & O'Neill, K. M. (1990?). *The Natural History of North American Beewolves*. Ithaca, NY: Cornell Univ. Press.
- Fain, A., & Gerson, U. (1990). Notes on two astigmatic mites (Acari) living in beehives in Thailand. *Acarologia*, 31(4), 381-384.
- Fairey, D. T., Lefkovitch, L. P., Lieverse, J. A., & Siemens, B. (1988). Materials for leafcutting bee (*Megachile rotundata* F.) shelters in north west Canada. *Zeitschrift fur Angewandte Entomologie*, 106, 119-122.
- Falcao, T. M. M.A., & Contel, E. P. B. (1990). Genetic variability in natural populations of Brazilian social bees. I. Isozyme patterns and polymorphism for esterases and total protein. *Rev. Bras. Genet.*, 13(4), 731-754.
- Fan, J. G. (1990). A study on Chinese *Halictus* (*Halictus*) with a description of three new species (Hymenoptera: Halictidae). *Acta Zootaxonomica Sin.*, 15(1), 92-97.
- Fan, J. G., & Wu, Y. R. (1991). Three new species of *Lasioglossum* from China (Hymenoptera: Halictidae). *Acta Entomol. Sin.*, 34(1), 89-93.
- Farina, W. M., & Nunez, J. A. (1991). Trophallaxis in the honeybee, *Apis mellifera* (L.) as related to the profitability of food sources. *Animal Behaviour*, 42(3), 389-395.
- Fernandes-da-Silva, P. G., & Zucoloto, F. S. (1990). A semi-artificial diet for *Scaptotrigona depilis* Moure (Hymenoptera: Apidae). *J. Apic. Res.*, 29(4), 233-235.
- Ferreira, L. M., de Albuquerque, C. M. R., & Hadj-Idris, A. E. Q. (1989). Aspectos bionómicos em *Melipona scutellaris* (Hymenoptera, Apidae, Meliponinae): postura de operárias e aprovisionamento. *Rev. Bras. Entomol.*, 33(2), 217-224.
- Fisher, R. M. (1987). Queen-worker conflict and social parasitism in bumble bees (Hymenoptera: Apidae). *Animal Behav.*, 35(4), 1026-1036.
- Fisher, R. M. (1987). Temporal dynamics of facultative social parasitism in bumble bees (Hymenoptera: Apidae). *Animal Behav.*, 35(6), 1628-1636.
- Fisher, R. M., & Pomeroy, N. (1989). Incipient colony manipulation, nosema incidence and colony productivity of the bumble bee *Bombus terrestris* (Hymenoptera: Apidae). *J. Kansas Entomol. Soc.*, 62(4), 581-589.
- Fisher, R. M., & Pomeroy, N. (1990). Sex discrimination and infanticide by queens of the bumble bee *Bombus terrestris* (Hymenoptera: Apidae). *Anim. Behav.*, 39 (4), 801-802.
- Francke, W., Krohn, S., & Tengo, J. (1991). Identification of new sesquiterpenoids in cephalic secretion of cuckoo bee *Nomada lathburiana* (Apoidea: Anthophoridae). *J. Chem. Ecol.*, 17(3), 557-566.
- Frankie, G. W., Vinson, S. B., Newstrom, L. E., Barthell, J. F., Haber, W. A., & Frankie, J. K. (1991). Plant phenology, pollination ecology, pollinator behavior and conservation of pollinators in neotropical dry forest. In: K. S. Bawa, & M. Hadley, *Man and the Biosphere Series*, Vol. 7. *Reproductive Ecology of Tropical Forest Plants*; Meeting, Bangi, Malaya, June 8-12, 1987. (pp. 37-48). USA, England, France: Parthenon Publishing Group, Inc.
- Frankie, G. W., Vinson, S. B., & Williams, H. (1989). Ecological and evolutionary sorting of 12 sympatric species of *Centris* bees in Costa Rican dry forest. In: J. H. Bock, & Y. B. Linhart, *The Evolutionary Ecology of Plants*. (pp. 536-549). USA & London: Westview Press.
- Fritz, A. L. (1990). Deceit pollination of *Orchis spitzelii* (Orchidaceae) on the island of Gotland in the Baltic: a suboptimal system. *Nordic J. Botany*, 9(6), 577-587.
- Fritz, M., & Toro, H. (1989). The type specimens of *Coelioxys* (Hymenoptera: Megachilidae) described by E. L. Holmberg. *Rev. Chil. Entomol.*, 17, 73-78.
- Frohlich, D. R. (1990). Substrate specificity of esterases in a solitary bee, *Megachile rotundata* (Hymenoptera: Megachilidae): variability in sex, age and life stage. *Biochem. Syst. Ecol.*, 18(7/8), 530-547.
- Frohlich, D. R., Brindley, W. A., Burris, T. E., & Youssef, N. N. (1990). Esterase isozymes in a solitary bee, *Megachile rotundata* (Fab.): characterization, developmental multiplicity, and adult variability. *Biochem. Genet.*, 28(7/8), 347-358.
- Gasparian, A. O. (1990). The bee feels the earthquake. *Biol. Zh. Arm.*, 43(2), 127-129.
- Geiser, F. (1988). *Wildbienen: Wehrhafte Blumenkinder*. Hannover, Germany: Landbuch-Verlag GMBH, pp.1-135.
- Gerlach, G., & Schill, R. (1989). Fragrance analyses, an aid to taxonomic relationships of the genus *Coryanthes* (Orchidaceae). *Plant Syst. Evol.*, 168(3/4), 159-165.

- Gess, F. W., & Gess, S. K. (1990). Solitary bees: the little known and unsung thousands. *Naturalist* (Port Elizabeth), 34(1), 23-32.
- Giblin-Davis, R. M., Norden, B., Batra, S. W. T., & Eickwort, G. C. (1990). Commensal nematodes in the glands, genitalia, and brood cells of bees (Apoidea). *J. Nematology*, 22(2), 150-161.
- Gilbert, F. S., Haines, N., & Dickson, K. (1991). Empty flowers. *Funct. Ecol.*, 5(1), 29-39.
- Gilliam, M., Buchmann, S. L., Lorenz, B. J., & Schmalzel, R. J. (1990). Bacteria belonging to the genus *Bacillus* associated with three species of solitary bees. *Apidologie*, 21(2), 99-105.
- Gilliam, M., Roubik, D. W., & Lorenz, B. J. (1990). Microorganisms associated with pollen, honey, and brood provisions in the nest of a stingless bee, *Melipona fasciata*. *Apidologie*, 21(2), 89-97.
- Goerzen, D. W. (1991). Microflora associated with the alfalfa leafcutting bee, *Megachile rotundata*. *Apidologie*, 22(5), 553-561.
- Goerzen, D. W., Erlandson, M. A., & Moore, K. C. (1990). Effect of two insect viruses and two entomopathogenetic fungi on larval and pupal development in the alfalfa leafcutting bee *Megachile rotundata* Fab. (Hymenoptera: Megachilidae). *Can. Entomol.*, 122 (9-10), 1039-1040.
- Goerzen, D. W., Erlandson, M. A., & Bissett, J. (1990). Occurrence of chalkbrood caused by *Ascospaera aggregata* Skou in a native leafcutting bee *Megachile relativa* Cresson (Hymenoptera: Megachilidae) in Saskatchewan Canada. *Can. Entomol.*, 122(11-12), 1269-1270.
- Gogala, A. (1990-1991). Halictine - the life of wild bees. (Slovenian). *Proteus*, (83-89).
- Gogala, A. (1990). Methods of transporting pollen in bees (Slovenian). *Proteus*, 52(8), 337-339.
- Gonzalez, J. M., & Gaiani, M. A. (1989). New species of *Eufriesia* (Hymenoptera: Apidae) from Venezuela. *Rev. Biol. Trop.*, 37(2), 149-152.
- Gonzalez, J. M. (1991). El genero *Eulaema* (Apidae: Euglossini) en Venezuela, su distribucion e interacciones con diversas familias botanicas. (Abstract). Congreso Venezolano de Entomologia, Merida, del 1 al 4 de Julio de 1991, p.203.
- Gonzalez, J. M., & Gaiani, M. A. (1991). *Euglossini* del Cerro Aracamuni territorio federal amazonas, Venezuela (Hymenoptera: Apidae). *Acta Terramaris*, (3), 19-21.
- Gosek, J., Ruszowski, A., Bilinski, M., & Kaczmarska, K. (1988). The percentage of females in the progeny of the lucerne leafcutter bee (*Megachile rotundata*) in relation to the percentage of females in the parent generation and to the breeding line and population density. *Pszczelnicze Zeszyty Naukowe*, 32, 99-110.
- Gottsberger, G., & Silberbauer-Gottsberger, I. (1988). Evolution of flower structures and pollination in neotropical Cassiinae (Caesalpiniaceae) species. *Phyton*, Austria, 28(2), 293-320.
- Grewal, G. S., Singh, G., & Kandoria, J. L. (1990). Insect pollinators of pigeonpea (*Cajanus cajan*) around Ludhiana. *Indian J. Agric. Sci.*, 60(3), 227-228.
- Grieve, H., & Surholt, B. (1990). Dependence of fructose-bis-phosphate from flight muscles of the bumblebee (*Bombus terrestris* L.) on calcium ions. *Comp. Biochem. Physiol. B Comp. Biochem.*, 97(1), 197-200.
- Griswold, T. L. (1991). A review of the genus *Microthurge* (Hymenoptera: Megachilidae). *Pan-Pac. Entomol.*, 67(2), 115-118.
- Gross, H. R., & Carpenter, J. E. (1991). Role of the fall armyworm (Lepidoptera: Noctuidae) pheromone and other factors in the capture of bumblebees (Hymenoptera: Apidae) by universal moth traps. *Environ. Entomol.*, 20(1), 377-381.
- Gupta, M., & Kasana, V. K. (1990). Protection of bees from pesticides I. Field test of repellents on the foraging behavior of *Apis florea* F. *Indian J. Anim. Res.*, 24(2), 101-109.
- Gupta, R. K. (1991). Descriptions of the male genitalia of some Indian megachilid bees (Insects: Hymenoptera: Apoidea). *Entomol. Abh.*, 54(1-4), 97-104.
- Gupta, R. K. (1990). A new species of genus *Anthocopa* Lepeletier and Serville (Hymenoptera: Apoidea: Megachilidae) from Orissa India. *J. Bombay Nat. Hist. Soc.*, 86(3), 419-422.
- Gupta, R. K. (1990). A new species of genus *Creightonella* Cockerell (Hymenoptera: Apoidea: Megachilidae) from central India. *J. Bombay Nat. Hist. Soc.*, 86(3), 416-419.
- Gupta, R. K. (1990). On a new subgenus *Neoashmeadiella* new subgenus and two new species of genus *Ashmeadiella* Cock. from India (Insecta: Hymenoptera: Apoidea: Megachilidae). *Reichenbachia*, 28(1-19), 55-58.

- Haeseler, V. (1990). The wild bees of the east Frisian Island of Norderney (Hymenoptera: Apoidea). *Faun-Oekol. Mitt.*, 6(3-4), 125-146.
- Halada, J. (1989). The presence of the bee *Andrena vaga* in sandstone areas of Vlkov. (In Czech.). *Ziva*, 37(5), 225-226.
- Hallmen, M. (1990). Eine kolonie der Wildbienenart *Andrena vaga* (Panzer) im Naturschutzgebiet 'am Berger hang' (Hymenoptera, Apoidea, Andrenidae). *Hess. Faun. Briefe*, 10(1), 1-3.
- Hallmen, M. (1990). Einige Aspekte zur Erstansiedelung von Hummeln der Gattung *Bombus* (Hymenoptera: Apidae) im Freiland in künstlichen Nestern. *Nachr. Entomol. ver Apollo*, 11(1), 49-59.
- Hallmen, M., & Hallmen, K. (1989). Active species protection through uncleared house walls as exemplified in the solitary bee *Osmia cornuta* Latr. (Hymenoptera: Megachilidae). *Jahresber. Wetterausischen. Ges Gesamte. Naturkd. Hanau.*, 140-141, 45-52.
- Hamilton, W. W. (1988). Temporal changes in nectar availability and *Bombus appositus* (Hymenoptera: Apidae) foraging profits. *Southwest Nat.*, 33(2), 219-227.
- Harder, L. D. (1990). Behavioral responses by bumble bees to variation in pollen availability. *Oecologia*, 85(1), 41-47.
- Harder, L. D. (1990). Pollen removal by bumble bees and its implications for pollen dispersal. *Ecol. Publ. Ecol. Soc. Am.*, 71(3), 1110-1125.
- Hartfelder, K. H. (1987). Rates of juvenile hormone synthesis control caste differentiation in the stingless bee *Scaptigona postica* Depilis. *Roux's Archives of Developmental Biology*, 196, 522-526.
- Hartfelder, K. (1990). Regulatory steps in caste development of eusocial bees. In: W. Engels, *Social Insects: an Evolutionary Approach to Castes and Reproduction*. (pp. 245-264). New York: Springer-Verlag.
- He, W., & Wu, Y. (1990). Two new species of bees from Yunnan China (Hymenoptera: Apoidea). *Sinzoologica*, (7), 217-220.
- Hedstrom, I. (1989). Pollen carriers and fruit development of *Psidium guajava* L. (Myrtaceae) in the neotropical region. *Rev. Biol. Trop.*, 36(2B), 551-553.
- Heemert, C. v., Ruijter, A. d., Eijnde, J. v. den, & Steen, J. v. der. (1990). Year-round production of bumble bee colonies for crop pollination. *Bee World*, 71(2), 54-56.
- Heras, C., & Gayubo, S. F. (1989). Apido fauna de la Provincia de Zamora 2. Megachilidae (Hymenoptera, Apoidea). *Bol. Asoc. Esp. Entomol.*, 13, 237-249.
- Heras, C., & Gayubo, S. F. (1990). Contribution al conocimiento de los Apoideos de la Provincia de Zamora. 1. Anthophoridae (Hymenoptera: Apoidea). *Eos-Rev. Esp. Entomol.*, 65(1), 61-71.
- Hermosa de Mendoza, M., Hermosa de Mendoza, J., Puerta, F., Asenio, E., Bustos, M., Padilla, F., & Pellin, P. (1989). Ascospaeriosis of the parasitic bee, *Coelioxys rufocaudata* by *Ascospaera aggregata*. *J. Apic. Res.*, 28(2), 61-65.
- Herrera, C. M. (1990). Bumble bees feeding on non-plant food sources. *Bee World*, 71(2), 67-69.
- Hogendoorn, K., & Velthuis, H. H. W. (1988). Influence of multiple mating on kin recognition by worker honeybees. *Naturwissenschaften*, 75, 412-413.
- Horvitz, C. C., & Schemske, D. W. (1990). Spatiotemporal variation in insect mutualists of a neotropical herb. *Ecol. Publ. Ecol. Soc. Am.*, 71(3), 1085-1097.
- Houston, T. F. (1990). Descriptions of new paracolletine bees associated with flowers of *Eremophila* (Hymenoptera: Colletidae). *Rec. West. Aust. Mus.*, 14(4), 583-622.
- Houston, T. F. (1991). Two new and unusual species of the bee genus *Leioproctus* Smith (Hymenoptera: Colletidae), with notes on their behavior. *Rec. West Aust. Mus.*, 15(1), 83-96.
- Ikudome, S., & Yamane, S. (1990). The distribution of megachilid bees in the Ryukyu Islands, Japan (Hymenoptera, Apoidea). *Bull. Inst. Minami-Kyushu Reg. Sci.*, (6), 73-93.
- Inoue, T. (1990). A trip in Yucatan, Mexico - Meliponiculture of the Maya. *Honeybee Science*, 11(2), 49-58.
- Inoue, T., Adri, & Salmah, S. (1990). Nest site selection and reproductive ecology of the Asian honey bee, *Apis cerana indica* in central Sumatra. In: S. F. Sakagami, R. I. Ohgushi, & D. W. Roubik, *Natural History of Social Wasps and Bees in Equatorial Sumatra*. (pp. 219-232). Sapporo, Japan: Hokkaido University Press.
- Inoue, T., Salmah, S., Sakagami, S. F., Yamane, S., & Kato, M. (1990). An analysis of antophilous insects in Central Sumatra. In: S. F. Sakagami, R. I. Ohgushi, & D. W. Roubik, *Natural History of Social Wasps and Bees in Equatorial Sumatra*. (pp. 175-200). Sapporo, Japan: Hokkaido University Press.

- Iuga, V. G. (1989). La structure de la region gnathale n'etaye pas l'origine Sphecoideenne des Apoides. *Trav. Mus. Hist. Nat. 'Grigore Antipa'*, 30, 41-56.
- Jacob-Remacle, A., & Jacob, J. P. (1990). Interet faunistique des sablières de Lorraine Belge: l'exemple des hymenopteres apoides solitaires. *Notes Fauniques Gembloux*, 21, 13-22.
- Jacob-Remacle, A. (1990). Etude descriptive des nids d'*Osmia cornuta* (Latreille, 1805) et d'*Osmia rufa* (Linne, 1758) (Hymenoptera: Apoidea: Megachilidae). *Notes Fauniques de Gembloux*, (22), 49-64.
- Jaeschke, G. (1990). Die Zusammensetzung der Hummel - und Schmarotzerhummelfauna in Berlin-Pankow anhand von Totfunden (Hymenoptera: Apoidea: *Bombus* et *Psithyrus*). *Novius*, (10), 211-212.
- Jahns, T. R., & Joliff, G. D. (1991). Survival rate and reproductive success of *Osmia lignaria propinqua* Cresson (Hymenoptera: Megachilidae) in caged meadowfoam, *Limnanthes alba* Benth (Limnanthaceae). *J. Kansas Entomol. Soc.*, 64(1), 95-106.
- Janzon, L. A., Svensson, B. G., & Erlandsson, S. (1991). Catalogus insectorum sueciae (Hymenoptera: Apoidea) 3. Megachilidae, Anthophoridae and Apidae. *Entomol. Tidskr.*, 112(3), 93-99.
- Javier Ortiz, F. (1989). *Xylocopa cantabrita* Lepeletier, 1841 y *Xylocopa uclesiensis* Perez, 1901 en el sur de la Peninsula iberica (Hym. Anthophoridae). *Bol. Asco. Esp. Entomol.*, 13, 452.
- Jaycox, E. R., & Karpowicz, J. (1990). A beekeeping project in the Yemen Arab Republic. In: R. B. Serjeant, & R. L. Bidwell, *Arabian Studies*. (pp. 1-10). Great Britain: Cambridge University Press.
- Jelinski, M. (1988). Mass invasion of pyralids in a bumble bee nest in a bird's nest. (In Polish). *Wszechswiat*, 89(12), 289.
- Jennersten, O., & Kwak, M. M. (1991). Competition for bumblebee visitation between *Melampyrum pratense* and *Viscaria vulgaris* with healthy and *Ustilago*-infected flowers. *Oecologia*, 86(1), 88-98.
- Jennersten, O., & Morse, D. H. (1991). The quality of pollination by diurnal and nocturnal insects visiting common milkweed *Asclepias syriaca*. *Am. Midl. Nat.*, 125(1), 18-28.
- Johnson, M. D. (1990). Female size and fecundity in the small carpenter bee, *Ceratina calcarata* (Robertson) (Hymenoptera: Anthophoridae). *J. Kansas Entomol. Soc.*, 63(3), 414-419.
- Joos, B., Young, P. A., & Casey, T. M. (1991). Wingstroke frequency of foraging and hovering bumblebees in relation to morphology and temperature. *Physiol. Entomol.*, 16(2), 191-200.
- Jyothi, P. V., Atluri, J. B., & Reddi, C. S. (1990). Pollination ecology of *Moringa oleifera* Moringaceae. *Proc. Indian Acad. Sci. Plant Sci.*, 100(1), 33-42.
- Kaitala, V., Smith, B. H., & Getz, W. M. (1990). Nesting strategies of primitively eusocial bees: models of nest usurpation during the solitary state of the nesting cycle. *J. Theor. Biol.*, 144(4), 445-471.
- Katayama, E. (1989). Comparative studies on the egg-laying habits of some Japanese species of bumblebees (Hymenoptera, Apidae). *Occasional Publications, Entom. Soc. Jpn.*, (2), 1-161.
- Katayama, E., Ochiai, H., & Takamizawa, K. (1990). Supplementary notes on nests of some Japanese bumblebees. 2. *Bombus ussurensis*. *Jpn. J. Entomol.*, 58(2), 335-346.
- Kato, M. (1988). Bumblebee visits to *Impatiens* spp.: Pattern and efficiency. *Oecologia*, 76 (3), 364-370.
- Kawashima, K. (1989). Effect of chitin synthesis inhibitors on the horn faced bee, *Osmia cornifrons* Radoszowski. *J. Soc. Plant Protect. North Jpn.*, (40), 171-173.
- Keighery, G. J. (1991). Pollination of *Hibbertia conspicua* Dilleniaceae. *West. Aust. Nat.*, 18(6), 163-165.
- Kelber, A., & Zeil, J. (1990). A robust procedure for visual stabilization of hovering flight position in guard bees of *Trigona angustula* (Apidae: Meliponinae). *J. Comp. Physiol. A Sens. Neural Behav. Physiol.*, 167(4), 569-578.
- Kerr, W. E. (1987). Sex determination in bees. 24. Critical appraisal of sex determination in bees. (1). *Honeybee Sci.*, 8(4), 176-182.
- Kerr, W. E. (1990). Why are workers in social Hymenoptera not males? *Rev. Bras. Genet.*, 13(1), 133-136.
- Kerr, W. E., & Da Cunha, R. A. (1990). Sex determinintation in bees. 26. Masculinism of workers in the Apidae. *Rev. Bras. Genet.*, 13(3), 479-489.
- Kerr, W. E., Monteiro, S. G., & Kerr, H. A. S. (1988). Sex determination in bees. XXV. Adaptive value of the XO1 gene in its origin. *Rev. Bras. Genet.*, 11(2), 469-473.
- Kevan, P. G. (1990). How large bees *Bombus* and *Xylocopa* (Apoidea: Hymenoptera) forage on trees optimally and patterns of movement in temperate and tropical climates. *Ethol. Ecol. Evol.*, 2(3), 233-242.

- Kevan, P. G., Eisikowitch, D., Ambrose, J. D., & Kemp, J. R. (1990). Cryptic dioecy and insect pollination in *Rosa setigera* Michx. (Rosaceae), a rare plant of Carolinian Canada. *Biol. J. Linn. Soc.*, 40, 229-243.
- Kifune, T. (1991). Description of four new species of the genus *Halictoxenos* Strepsiptera Stylopidae from Japan. *Jpn. J. Entomol.*, 59(2), 367-374.
- Kifune, T. (1991). Two new species of the genus *Stylops* (Strepsiptera: Stylopidae) with stylopized records of the andrenid bees in Japan. (Studies on the Japanese Strepsiptera XIV). *Japan J. Entomol.*, 59(1), 155-163.
- Kifune, T., & Maeta, Y. (1990). Ten new species of the genus *Stylops* (Strepsiptera: Stylopidae) parasitic on the genus *Andrena* (Hymenoptera: Andrenidae) of Japan studies on the Japanese Strepsiptera XIII. *Esakia*, (Spec. Issue 1), 97-110.
- Kim, M. L., & Kim, C. W. (1989). Numerical phenetic analysis of the genus *Andrena* from Korea (Hymenoptera: Andrenidae). *Entomol. Res. Bull.*, 15, 25-33.
- Klinkhamer, P. G. L., & de Jong, T. J. (1990). Effects of plant size, plant density and sex differential nectar reward on pollinator visitation in the protandrous *Echium vulgare* (Boraginaceae). *Oikos*, 57(3), 399-405.
- Klinkhamer, P. G. L., de Jong, T. J., & Bruyn, G. J. de. (1989). Plant size and polinator visitiation in *Cynoglossum officinale*. *Oikos*, 54(2), 201-204.
- Koeniger, G., Koeniger, N., Mardan, M., Otis, G., & Wongsiri, S. (1991). Comparative anatomy of male genital organs in the genus *Apis*. *Apidologie*, 22(5), 539-552.
- Koeniger, G., Koeniger, N., Mardan, M., Punchihewa, R. W. K., & Otis, G. W. (1990). Numbers of spermatozoa in queens and drones indicate multiple mating of queens in *Apis andreniformis* and *Apis dorsata*, *Apidologie*, 21(4), 281-286.
- Kohl, A. (1989). Untersuchung von eingetragenem Pollen bei in künstlichen Nestern gehaltenen Hummelarten (Hymenoptera, Apoidea) und Rekonstruktion der besuchten Phytozonenosen im Jahresverlauf. *Ges. Oekol. Verh.*, 17, 713-718.
- Kosior, A., & Nosek, A. (1987). Species composition and number of bumblebees *Bombus* Latr. in the areas influenced by the emission from non-ferrous metal works in the Silesian Upland. *Studia Natura*, (31), 81-99.
- Krannitz, P. G., & Maun, M. A. (1991). Insect visitors to the Guelder Rose, *Viburnum opulus* var. *opulus* (Caprifoliaceae), in London, Ontario. *Can. Field. Nat.*, 105(1), 13-17.
- Kratochwil, A. (1989). Community structure of flower-visiting insects in different grassland types in southwestern Germany (Hymenoptera, Apoidea, Lepidoptera, Diptera). *Spixiana* (Muench), 12(3), 289-302.
- Kratochwil, A., & Klatt, M. (1989). Apoide Hymenopteren an Ruderalfstellen der Stadt Freiburg I. (BRD) - Sudmediterrane Faunenelemente an Standorten kleinraumig hoher Persistenz. *Zool. Jahrb. Abt. Syst. Oekol. Geogr. Tiere.*, 116(4), 379-389.
- Kreisch, W., & Schick, B. (1989). Bevorzugt die Wildbiene *Anthidium manicatum* einige wenige Fulterpflanzen? *Mitt. Pollichia*, 76 (171-183).
- Kress, A. (1988). Observations on *Kalmia latifolia* flowers (Ericaceae). *Phytologia*, 65(4), 249-284.
- Kukuk, P. F. (1990). Diploid males in a primitively eusocial bee, *Lasioglossum (Dialictus) zephyrum* (Hymenoptera: Halictidae). *Evolution*, 44(6), 1522-1528.
- Kukuk, P. F., & Crozier, R. H. (1990). Trophallaxis in a communal halictine bee *Lasioglossum (Chilalictus) erythrurum*. *Proc. Natl. Acad. Sci. USA*, 87(14), 5401-5404.
- Kukuk, P. F., & May, B. (1991). Colony dynamics in a primitively eusocial halictine bee *Lasioglossum (Dialictus) zephyrum* (Hymenoptera: Halictidae). *Insectes Sociaux*, 38(2), 171-188.
- Kumar, J., Rao, K. V. K.K., Gupta, P. R., & Dogra, G. S. (1989). Temporal distribution of bees (Hymenoptera: Apoidea) on spring flowering crops in midhills of Himachal Pradesh, India. *Indian Bee J.*, 51(2), 55-58.
- Kumar, V. (1989). A new species of the genus *Osmia* Panzer (Hymenoptera: Apoidea: Megachilidae) from India. *J. Entomol. Res.*, 13(1-2), 137-139.
- Kumar, V. (1989). A new species of the genus *Coelioxys* Latreille (Hymenoptera: Apoidea: Megachilidae) from India. *J. Entomol. Res.*, 13(1-2), 128-130.
- Kumar, V. (1990). A new species of *Megachile* Latreille (Hymenoptera: Apoidea: Megachilidae) from India. *J. Bombay Nat. Hist. Soc.*, 87(2), 266-268.

- Kumar, V., & Tiwari, V. K. (1990). A new species of genus *Parevaspis ritsema* (Hymenoptera: Apoidea: Megachilidae: Anthidinae) from India. *J. Bombay Nat. Hist. Soc.*, 87(1), 122-124.
- Kwak, M. M., & Jennersten, O. (1991). Bumblebee visitation and seedset in *Melampyrum pratense* and *Viscaria vulgaris*: heterospecificity pollen and pollen limitation. *Oecologia*, 86(1), 99-104.
- Labougle, J. M. (1990). *Bombus* of Mexico and Central America (Hymenoptera, Apidae). *Univ. Kansas Sci. Bull.*, 54(3), 35-73.
- Lack, A. J., & Kevan, P. G. (1987). The reproductive biology of a distylous tree, *Sarcocapnos celebica* (Oxalidaceae), in Sulawesi, Indonesia. *Bot. J. Linnean Soc.*, 95(1), 1-8.
- Laroca, S. (1990). Sobre visitas de Abelhas silvestres (Apoidea) as flores de *Brachiaria humidicola* (Gramineae), em porto Velho (Rondonia, Brasil). *An. Soc. Entomol. Bras.*, 19(2), 481-484.
- Laroca, S., & Almeida, M. C. D. (1989). Coexistence between stingless bees and ants nest of *Paratrigona myrmecophila* Apidae constructed in a nest of *Camponotus senex* Formicidae. *Rev. Bras. Zool.*, 6(4), 671-680.
- Larson, K. S., & Larson, R. J. (1990). Lure of the locks: showiest ladies-tresses orchids. *Spiranthes romanzoffiana*, affect bumblebee, *Bombus* spp., foraging behavior. *Can. Field Nat.*, 104(4), 519-525.
- Larsson, F. K. (1990). Female body size relationships with fecundity and egg size in two solitary species of fossorial Hymenoptera (Colletidae and Sphecidae). *Entomol. Gen.*, 15(3), 167-171.
- Larsson, F. K. (1989). Insect mating patterns explained by microclimatic variables. *J. Thermal Biol.*, 14(3), 155-157.
- Larsson, F. K. (1991). Some like it cool, some like it hot - a comparative study of male mate searching tactics in two species of Hymenoptera (Colletidae and Specidae). *J. Therm. Biol.*, 16(1), 45-51.
- Larsson, F. (1989). Matings patterns in six insect species: effects of weather and population density. *Acta Univ. Ups.*, Comprehensive Summaries of Uppsala Dissertations from the Faculty of Science 234., 33pp.
- Lehrer, M. (1990). How bees use peripheral eye regions to localize a frontally positioned target. *J. Comp. Physiol. A Sens. Neural Behav. Physiol.*, 167(2), 173-186.
- Leigh Jr., E. G. (1991). Genes, bees, and ecosystems: the evolution of a common interest among individuals. *Trends Ecol. Evol.*, 6(8), 257-262.
- Li, S. W., Meng, Y. P., Chang, J. T., & Li, J. H. (1987). A comparative study of esterase isozymes in Hymenoptera. *Acta Entomologica Sinica*, 3(30), 266-270.
- Lin, T. C. (1990). The abundance and diversity of Hymenopterans in Ulu Kinchin Pahang Malaysia. *Malay. Nat.*, 43(4), 278-281.
- Lobreaux-Callen, D. (1989). Les Malpighiaceae et leurs polliniseurs. Coadaptation ou coévolution. *Bulletin du Museum National D'Histoire Naturelle*, 11(1), 79-94.
- Lobreaux-Callen, D., Le Thomas, A., Darchen, B., & Darchen, R. (1990). Quelques facteurs déterminant le comportement de butinage d'*Hypotrigona pothieri* (Trigonini) dans la végétation de côte-d'ivoire. *Apidologie*, 21 (1), 69-83.
- Lunau, K. (1990). Colour saturation triggers innate reactions to flower signals: flower dummy experiments with bumblebees. *J. Comp. Physiol. A Sens. Neural Behav. Physiol.*, 166(6), 827- 834.
- Maa, T. C. (1987). An Enumeration of the *Xylocopa* bees of Fujian, China. *Wuyi Sci. J.*, 7, 205-210.
- MacFarlane, R. P., & Griffin, R. P. (1990). New Zealand distribution and seasonal incidence of the nematode, *Sphaerularia bombi* Dufour, a parasite of bumble bees. *N. Z. J. Zool.*, 17(2), 191-199.
- Machado, M. F. P.S., & Contel, E. P. B. (1991). Glycerol-3-phosphate dehydrogenase (G-3-PDH; EC 1.1.1.8) variation in Brazilian stingless bees and in wasp species. *Biochem. Genet.*, 29(5/6), 255-260.
- Machado, V. L. L., Giannotti, E., & de Oliveira, R. M. (1988). Insects visiting *Terpanax papyriferus* during its flowering period. *Rev. Bras. Biol.*, 48(3), 537-544.
- Macior, L. W. (1990). Pollination ecology of *Pedicularis punctata* Decne. Scrophulariceae in the Kashmir Himalaya. *Plant Species Biol.*, 5(2), 215-224.
- Maeta, Y., Sasaki, Y., & Fujimoto, G. (1988). *Andrena postomias* of the Gakuonji temple in hyogo prefecture. *Insectarium*, 25(2), 50-57.
- Maeta, Y., & Sugiura, N. (1990). Decision-making in a mason bee, *Osmia cornifrons* (Radoszkowski) (Hymenoptera, Megachilidae): does the mother bee fertilize her eggs depending on their sizes? *Jpn. J. Entomol.*, 58(1), 197-203.

- Maki, D. L., Moffett, J. O., & Petersen, H. D. (1990). X- radiography effects on leafcutting bee, *Megachile rotundata*, body size and survival at emergence. Southwest Entomol., 15(2), 147-150.
- Manino, A., & Patetta, A. (1989). A comparison of intestinal microvilli in bumble bees. Apicoltore Moderno, 80(3), 129-136.
- Marikovskaya, T. P. (1990). New data on the biology of bees (Hymenoptera, Apoidea) of mountains and submontane plains of South Kazakhstan. Tr. Inst. Zool. Akad. Nauk. Kaz. SSR, 45, 149-158.
- Marikovskaya, T. P., & Shcherbakova, T. I. (1990). Biology of *Osmia rufa* L. and *Osmia cornuta* Latr. (Hymenoptera, Megachilidae) inhabiting artificial nests in the foothills of Zailiisk Alatau. Tr. Inst. Zool. Akad. Nauk. Kaz. SSR, 45, 135-148.
- Mayer, D. F., Lunden, J. D., & Miliczky, E. R. (1990). Effects of fungicides on chalkbrood disease of alfalfa leafcutting bee. Appl. Agric. Res., 5(3), 223-226.
- Maynard, G. V. (1991). Revision of *Leioproctus protomorpha* Rayment (Hymenoptera: Colletidae) with a description of two new species. J. Aust. Entomol., 30(1), 67-75.
- McAuslane, H. J., Vinson, S. B., & Williams, H. J. (1990). Change in mandibular and mesosomal gland contents of male *Xylocopa micans* (Hymenoptera: Anthophoridae) associated with mating system. J. Chem. Ecol., 16(6), 1877-1885.
- McVetty, P. B. E., Pinnisch, R., & Scarth, R. (1989). The significance of floral characteristics in seed production of four summer rape cultivar a-lines with pol cytoplasm. Can. J. Plant Sci., 69(3), 915-918.
- Michener, C. D. (1990). Castes in Xylocopine bees. In: W. Engels, Social Insects: an Evolutionary Approach to Castes and Reproduction. (pp. 123-146). New York: Springer-Verlag.
- Michener, C. D. (1990). Reproduction and castes in social halictine bees. In: W. Engels, Social Insects: an Evolutionary Approach to Castes and Reproduction. (pp. 77-121). New York: Springer-Verlag.
- Michener, C. D., Brooks, R. W., & Pauly, A. (1990). Little-known Meganomiine bees with a key to the genera (Hymenoptera: Melittidae). Rev. Zool. Afr., 104(2), 135-140.
- Michener, C. D. (1990). Classification of the Apidae (Hymenoptera) Univ. Kansas Sci. Bull., 54(4), 75-164.
- Miliczky, E. (1991). Observations on the nesting biology of three species of panurgine bees (Hymenoptera: Andrenidae). J. Kansas Entomol. Soc., 64(1), 80-87.
- Miliczky, E. R., Mayer, D. F., & Lunden, J. D. (1990). Notes on the nesting biology of *Andrena (Melandrena) nivalis* Smith (Hymenoptera: Andrenidae). J. Kansas Entomol. Soc., 63(1), 166-174.
- Minckley, R. L., Buchmann, S. L., & Wcislo, W. T. (1991). Bioassay evidence for a sex attractant pheromone in the large carpenter bee *Xylocopa varipuncta* (Anthophoridae: Hymenoptera). J. Zool. (Lond.), 224(2), 285-292.
- Minckley, R. L., & Buchmann, S. L. (1990). Territory site selection of male *Xylocopa (Neoxylocopa) varipuncta* Patton (Hymenoptera: Anthophoridae). J. Kansas Entomol. Soc., 63(2), 329-339.
- Mingo, E., Gayubo, S. F., & Rueda, A. (1989). Contribucion al conocimiento de la familia Chrysididae de la provincia de Palencia (Hym.: Chrysididae). Eos-Rev. ESP Entomol., 65(2), 31-50.
- Monteiro, C. A., & Kerr, W. E. (1990). Experimental exchange of queens between colonies of *Melipona compressipes* (Apidae: Meliponini). Rev. Bras. Biol., 50(4), 975-982.
- Moritz, R. F. A., & Crewe, R. M. (1988). Air ventilation in nests of two African stingless bees *Trigona denoiti* and *Trigona gribodoi*. Experientia, 44(11/12), 1024-1027.
- Moritz, R. F. A., & Sakofski, F. (1991). The role of the queen in circadian rhythms of honeybees (*Apis mellifera* L.). Behav. Ecol. Sociobiol., 29(5), 361-367.
- Morris, R. K. A. (1991). *Hedychridium coriaceum* (Hymenoptera: Chrysididae) and other less common aculeate Hymenoptera from Mitcham Common Surrey England, UK. Entomol. Rec. J. Var., 103(5-6), 127-128.
- Moure, J. S. (1989). *Camargoia* new genus of Meliponinae from the neotropical region (Hymenoptera: Apoidea). Bol. Mus. Para Emilio Goeldi, 5(1), 71-78.
- Moure, J. S. (1988). A new species of *Frieseomelitta* from the western region of Amazonia (Hymenoptera: Apoidea). Acta Biol. Parana, 17(1-4), 141-146.
- Moure, J. S. (1989). New species of bees from the central region of Minas Gerais State Brazil (Hymenoptera: Apoidea). Acta Biol. Parana, 18(1-4), 115-127.

- Moure, J. S. (1989). *Sakagami affabria* new genus new species of (Meliponinae): Hymenoptera: Apidae) of Rondonia Brazil. Rev. Bras. Zool., 6(4), 681-684.
- Moure, J. S., Camargo, J. M. F., & Garcia, M. V. B. (1988). A new species of *Leurotrigona* (Hymenoptera: Apidae: Meliponinae). Bol. Mus. Para. Emilio Goeldi Ser. Zool., 4(2), 145-154.
- Murakami, K., & Itino, T. (1990). Foraging behavior of pollinator bees on chinese milk vetch *Astragalus sinicus* (Leguminosae) in relation to diurnal nectar-secreting pattern of the flowers. Honeybee Sci., 11(1), 11-16.
- Murcia, C. (1990). Effect of floral morphology and temperature on pollen receipt and removal in *Ipomoea trichocarpa*. Ecol. Publ. Ecol. Soc. Am., 71(3), 1098-1109.
- Naim, M., & Bisht, D. S. (1990). Fidelity and time spent on flowers by *Apis cerana* F. foragers. Indian J. Entomol., 51(2), 222-223.
- Narolsky, N. B., & Shcherbal, I. S. (1991). New data on Gasteruptidae (Hymenoptera, Evanoidae) - cleptoparasites of the leaf-cutter bee *Megachile rotundata*. Vestn. Zool., 1991(1), 22-24.
- Nates Parra, G., & Roubik, D. W. (1990). Sympatry among subspecies of *Melipona favosa* in Columbia and a taxonomic revision. J. Kansas Entomol. Soc., 63(1), 200-203.
- Naumann, I. D. (1990). The aculeate wasps and bees (Hymenoptera) of Norfolk and Philip Islands South Pacific Ocean. Aust. Entomol. Mag., 17(1), 17-28.
- Naumann, K. (1991). Grooming behaviors and translocation of queen mandibular gland pherome on worker honey bees (*Apis mellifera* L.). Apidologie, 22(5), 523-531.
- Naumann, K., Winston, M. L., Slessor, K. N., Prestwich, F. X., & Webster, F. X. (1991). Production and transmission of honey bee queen (*Apis mellifera* L.) mandibular gland pheromone. Behav. Ecol. Sociobiol., 29(5), 321-332.
- Nazarov, V. V., & Ivanov, S. P. (1990). Pollination of the mimetic species *Cephalanthera rubra* Z. Rich. and *Campanula taurica* Juz. by bees of the genus *Chelostoma* Latr. (Hymenoptera: Megachilidae) in the Crimea Ukrainian SSR USSR. Entomol. Obozr., 69(3), 534-537.
- Needham, G. R., Page Jr., R. E., Delfinado-Baker, M., & Bowman, C. E. (Eds.) (1988). Aricanized Honey Bees and Bee Mites. Chichester: Ellis Horwood Limited, 572pp.
- Neff, J. L., & Simpson, B. B. (1990). The role of phenology and reward structure in the pollination biology of wild sunflower (*Helianthus annus* L., Asteraceae). Israel J. Bot., 39, 197-216.
- Neff, J. L., & Simpson, B. B. (1991). Nest biology and mating behavior of *Megachile fortis* in Central Texas (Hymenoptera: Megachilidae). J. Kansas Entomol. Soc., 64 (3), 324-336.
- Negoro, H. (1990). Bumblebees from Toyama prefecture, Honshu, Japan. Bull. Toyama Sci. Mus., (13), 97-105.
- Nicolson, S. W. (1990). Osmoregulation in a nectar-feeding insect, the carpenter bee *Xylocopa capitata*: water excess and ion conservation. Physiol. Entomol., 15(4), 433-440.
- Nilsson, G. E. (1991). The wasp and bee fauna of the Rido archipelago in Lake Malaren, Sweeden (Hymenoptera, aculeata). Ent. Tidskr., 112, 79-92.
- Nobile, V. (1987). Contributo alla conoscenza degli Apoidei (Insecta, Hymenoptera) di Sicilia. 1. Generi *Habropoda* Smith, *Tetralonia* Spinola (Gruppo 'ruficornis F.') *Melecta* Latreille, *Eupavlovskia* Popov e *Thyreus* Panzer. Animalia (Catania), 14 (1-3), 73-89.
- O'Keefe, K. J., & Schwarz, M. P. (1990). Pheromones are implicated in reproductive differentiation in a primitively social bee. Naturwissenschaften, 77(2), 83-86.
- O'Neill, K. M., Evans, H. E., & Bjostad, L. B. (1991). Territorial behavior in males of three North American species of bumblebees (Hymenoptera: Apidae: *Bombus*). Can. J. Zool., 69(3), 604-613.
- Ornosa, C., & Gayubo, S. F. (1989). Notas sobre los Bombinae de la Sierra de Bejar (Hym., Apidae). Bol. Asoc. Esp. Entomol., 13, 173-182.
- Ortiz-Sanchez, F. J. (1991). New records to the knowlege of the genus *Nomoiodes* Schenck 1866 in the Iberian Peninsula (Hymenoptera: Halictidae). Eos-Rev. Esp. Entomol., 66(2), 157-160.
- Packer, L., Jessome, V., Lockerbie, C., & Sampson, B. (1989). The phenology and social biology of four sweat bees in a marginal environment: Cape Breton island. Can. J. Zool., 67(12), 2871- 2877.
- Packer, L., & Owen, R. E. (1990). Allozyme variation, linkage disequilibrium and diploid male production in a primitively social bee *Augochlorella striata* (Hymenoptera: Halictidae). Heredity, 65(Pt. 2), 241-248.

- Page Jr., R. E., & Kerr, W. E. (1990). The evolution of monandry and queen replacement in *Melipona* (Hymenoptera: Apidae). Rev. Bras. Genet., 13(2), 209-229.
- Pagliano, G. (1987). Prospetto Sistematico degli Apoidea Italiani. Annali Della Facolta di Scienze Agrarie Della Universita Delgi Studi di Torino, 38, 85-128.
- Pagliano, G., & Scaramozzino, P. (1989). Elenco dei gernerii di Hymenoptera del mondo. Memorie della Societa Entomologica Italiana, 68, 1-210.
- Parker, F. D., & Tepedino, V. J. (1990). Bee pollination of *Cuphea* (Lythraceae) species in greenhouse and field. Pan-Pac. Entomol., 66(1), 9-12.
- Parra Valencia, G. (1991). Bionomia de las abejas sin agujon (Apidae, Meliponinae) del occidente Colombiano. Cespedesia, 16-17 (57-58), 77-116.
- Patetta, A., & Manino, A. (1989). Convegno a Ferrara su artropodi sociali e presociali. Apiculture Moderno, 80(3), 121-125.
- Paulus, H. F., & Gack, C. (1990). Pollination of *Ophrys* (Orchidaceae) in cyprus. Plant Syst. Evol., 169(3-4), 177-207.
- Paulus, H. F., & Gack, C. (1990). Pollinators as prepollinating isolation factors: evolution and speciation in *Ophrys* (Orchidaceae). Isreal J. Bot., 39(1-2), 43-79.
- Pauly, A. (1990). Classification of African Nominae (Hymenoptera: Apoidea: Halictidae). K. Mus. Midden-Afr. Tervuren Belg. Ann. Zool. Wet., 261, 1-206.
- Pavlenko, G. V., & Naumenko, Z. S. (1987). Frateuria-like bacteria on (bumble) bees in the north-western RSFSR. Vestnik Leningradskogo Universiteta, Biologiya, 4 (24), 55-61.
- Pawlowski, T. (1989). The structure of wild bee (Hymenoptera: Apoidea) communities from farming areas of different field sizes. Acta Univ. Nicolai Copernici Biol., 33, 31-46.
- Pawlowski, T. (1989). The structure of wild bee (Hymenoptera: Apoidea) communities from habitats of field-pine forest ecotone. Acta Univ. Nicolai Copernici Biol., 33, 101-109.
- Pawlowski, T., & Pokorniecka, J. (1990). Observations on the structure of bumblebee communities Apoidea *Bombus* Latr. of the town-forest areas in Torun Basin North Poland. Acta Univ. Nicolai Copernici Biol., 37, 3-22.
- Paxton, R. J. (1991). Profile of a solitary bee: *Andrena fulva*. Bee World, 72(1), 11-18.
- Payette, A., & De Oliveira, D. (1990). Diversite et abondance des Apoides (Hymenoptera: Apoidea) dans l'agroecosysteme de Saint-Hyacinthe, Quebec. Nat. Can. (Que), 116(3), 155-165.
- Pekkarinen, A. (1988). Euro-Siberian element in the Fennoscandian bumble bee fauna (Hymenoptera, Apidae: *Bombus* and *Psithyrus*). In: V. V. Zlobin, Svyazi Entomofaun Severnoi Evropy I Sibiri. (pp. 118-125). Leningrad, USSR: Zoological Institute, USSR Academy of Sciences.
- Pesenko, Y. A. (1989). Bees of the tribe Nomiodini (Hymenoptera, Halictidae) of the transcaucasus and Asia minor, with a description of a new species of the genus *Nomioides* Schenck. Trudy Zool. Instit., Akad. Nauk SSSR, 188, 122-128.
- Pesenko, Y. A. (1988). Comparative analysis of the distribution of the genera *Halictus* Latreille S. Str. and *Lasioglossum* Curtis S. Str. (Hymenoptera, Halictidae) in the palaearctic. In: V. V. Zlobin, Svyazi Entomofaun Severnoi Evropy I Sibiri. (pp. 126-141). Leningrad, USSR: Zoological Institute, USSR Academy of Sciences.
- Pesenko, Y. A., Lelej, A. S., Radchenko, V. G., & Filatkin, G. N. (1990). The chinese wax bees *Apis cerana cerana* F. (Hymenoptera, Apidae) in the Soviet far east. Entomol. Rev., 69(3), 21-41.
- Pesenko, Y. A., & Situdikov, A. A. (1990). Classification and phylogenetic relationships between genera of the tribe Eucerini (Hymenoptera, Anthophoridae) with two submarginal cells. Entomol. Rev., 69(1), 88-104.
- Petit, J. (1990). Quelques donnees recentes sur la repartition et l'ethologie de *Macropis labiata* (Fabricius), abeille solitaire nouvelle pour la montagne Saint-Pierre. Nat. Mosana, 43(2), 29-39.
- Pintaudi, C. A., Stort, M. N. S., & Marin-Morales, M. A. (1990). Artifical and natural pollination of *Xylobium squalens* Lindl. Orchidaceae. Naturalia, 15, 67-80.
- Pleasants, J. M. (1989). Optimal foraging by nectarivores: a test of the marginal-value theorem. Am. Nat., 134(1), 51-71.
- Pleasants, J. M., & Zimmerman, M. (1990). The effect of inflorescence size on pollinator visitation of *Delphinium nelsonii* and *Aconitum columbianum*. Collect. Bot., 19, 21-40.

- Plowright, R. C., & Plowright, C. M. S. (1990). The laying of male eggs by bumble bee queens: an experimental reappraisal and a new hypothesis. *Can. J. Zool.*, 68(3), 493-497.
- Podbolotskaya, M. N. (1988). An analysis of the distribution of palaearctic bumble bees (Hymenoptera, Apidae, *Bombus* Latr.). In: V. V. Zlobin, *Svyazi Entomofaun Severnoi Evropy I Sibiri*. (pp. 142-147). Leningrad, USSR: Zoological Institute, USSR Academy of Sciences.
- Popova, L. M. (1990). Nesting habits of some species of anthophorid bees (Hymenoptera, Anthophoridae) in the middle Volga region. *Entomol. Rev.*, 69(5), 81-93.
- Popova, L. M. (1990). Nesting of several species of anthophorid bees (Hymenoptera) in the middle Volga area Russian SFSR USSR. *Entomol. Obozr.*, 69(1), 23-35.
- Possingham, H. P., Houston, A. I., & McNamara, J. M. (1990). Risk-averse foraging in bees: a comment on the model of Harder and Real. *Ecology*, 71(4), 1622-1624.
- Pouvreau, A. (1991). Morphology and histology of tarsal glands in bumble bees of the genera *Bombus*, *Ptychobombus*, and *Megabombus*. *Can. J. Zool.*, 69(4), 866-872.
- Priore, R. (1990). The Apoidea Hymenoptera collection of the Institute of Agarian Entomology in Portici Italy. X. *Celioxyx* Latr. *Megachilidae* *Xylocopa* Latr. *Ceratina* Latr. *Xylocopidae*. *Boll. Lab. Entomol. Agrar. Filippo. Silvestri.*, 46, 31-44.
- Radchenko, V. G., & Pesenko, Y. A. (1989). A key to bees of the genus *Dasyprocta* Latreille (Hymenoptera, Melittidae) of the European part of the USSR, with a designation of lectotypes. *Trudy Zool. Instit., Akademiya Nauk. SSSR*, 188, 114-121.
- Raju, A. J. S. (1989). Reproductive ecology of *Ocimum americanum* L. and *Ocimum basilicum* L. Lamiaceae in India. *Plant Species Biol.*, 4(2), 107-116.
- Raju, A. J. S., & Reddi, C. S. (1989). Pollination biology of *Anisomeles indica* and *Anisomeles malabarica* Lamiaceae. *Plant Species Biol.*, 4(2), 157-167.
- Ramalho, M. (1990). Foraging by stingless bees of the genus *Scaptotrigona* (Apidae, Meliponinae). *J. Apic. Res.*, 29(2), 61-67.
- Ramanujam, C. G. K., & Khatija, F. (1991). Melittopalynology of the agricultural tracts in Guntur district Andhra Pradesh India. *J. Indian Inst. Sci.*, 1(71), 25-34.
- Rank, G. H., & Rank, F. P. (1990). Pre-harvest diurnal temperature variations increase diapause development in the alfalfa leafcutting bee *Megachile rotundata* Fab. (Hymenoptera: Megachilidae). *J. Appl. Entomol.*, 110(3), 313-317.
- Rasmont, P., Barbier, Y., & Pauly, A. (1990). Faunistique comparee des Hymenopteres Apoides de deux terrils du Hainaut occidental. Notes Fauniques Gembloux, (21), 39-58.
- Ratnieks, F. L. W. (1991). Africanized bees natural selection for colonizing ability. In: M. Spivak, D. J. C. Fletcher, & M. D. Breed, *The "African" Honey Bee*. (pp. 119-136). Boulder, Colorado, USA; Oxford, England, UK: Westview Press.
- Read, P. E. C., Donovan, B. J., & Griffin, R. P. (1989). Use of bumblebees, *Bombus terrestris*, as pollinators of kiwifruit and lucerne in New Zealand. *N. Z. Entomol.*, 12, 19-23.
- Real, L. (1991). Animal choice behavior and the evolution of cognitive architecture. *Science (Washington, D.C.)*, 253(5023), 980-986.
- Real, L. A. (1990). Predator switching and the interpretation of animal choice behavior: the case for constrained optimization. *Nato Adv. Sci. Inst. Ser. Ser. G Ecol. Sci.*, 20, 1-21.
- Real, L. A., Ellner, S., & Harder, L. D. (1990). Short-term energy maximization and risk-aversion in bumble bees: a reply to Possingham et al. *Ecol.*, 71(4), 1625-1628.
- Reddy, T. B., & Aruna, C. (1990). Pollination ecology of *Alangium lamarckii* Alangiaceae. *Proc. Indian Acad. Sci. Plant Sci.*, 100(3), 195-204.
- Rego, M. M. C., & Albuquerque, P. M. C.D. (1990). Behavior of bees visiting murici trees *Byrsonima crassifolia* L. Kunth Malpighiaceae. *Bol. Mus. Para Emilio Goeldi Ser. Zool.*, 5(2), 179-184.
- Reimann, H. (1987). Die Bienen, Wespen und Ameisen (Hymenoptera, aculeata) der Naturschutzgebiete "Dunengebeit bei Neumuhlen" und "Vossberge" unter Berücksichtigung weiterer Binnendunenareale. Beiheft zur Schriftenreihe Naturschutz und Landschaftspflege in Niedersachsen: Hannover, (17), 79.
- Rembold, H. (1987). Die Kastenbildung bei der Honigbiene, *Apis mellifica* L., aus biochemischer Sicht. In: G. H. Schmidt, *Sozialpolymorphismus bei Insekten*. (pp. 350-403). Stuttgart: Wissenschaftliche Verlagsgesellschaft MBH.

- Renner, S. S. (1987). Reproduction and evolution in some genera of Neotropical Melastomataceae. *Mem. N.Y. Bot. Gard.*, 55, 143-152.
- Reyes, S. G. (1991). A review of the bee genus *Braunsapis* in Madagascar (Hymenoptera: Anthophoridae). *J. Afr. Zool.*, 105(2), 125-130.
- Reyes, S. G. (1991). Revision of the bee genus *Braunsapis* in the oriental region (Apoidea: Xylocopinae: Allodapini). *Univ. Kansas Sci. Bull.*, 54(6), 179-207.
- Reyes, S. G., & Michener, C. D. (1990). Observations on a parasitic allodapine bee and its hosts in Java and Malaysia (Hymenoptera: Anthophoridae: Allodapine). *Trop. Zool.*, 3(2), 139-149.
- Reyes, S. G., & Sakagami, S. F. (1990). A new socially parasitic *Braunsapis* from India with notes on the synonymy of *Braunsapis mixta* (Hymenoptera: Xylocopinae: Allodapini). *J. Kansas Entomol. Soc.*, 63(3), 458-461.
- Richards, K. W., & Krunic, M. D. (1990). Introduction of alfalfa leafcutter bees to pollinate alfalfa in Yugoslavia. *Entomologist*, 109(3).
- Richards, K. W., & Whitfield, G. H. (1988). Emergence and survival of leafcutter bees, *Megachile rotundata*, held at constant incubation temperatures (Hymenoptera: Megachilidae). *J. Apic. Res.*, 27(3), 197-204.
- Rico-Gray, V., & Thien, L. B. (1989). Some aspects of the reproductive biology of *Schomburgkia tibicinis* Batem. Orchidaceae in Yucatan Mexico. *Brenesia*, (28), 13-24.
- Riddick, E. W. (1990). *Andrena macra* Mitchell (Hymenoptera: Andrenidae) overwinter and delay spring emergence in Virginia. *Proc. Entomol. Soc. Wash.*, 92(4), 771-772.
- Riemann, H., & Melber, A. (1990). Hymenoptera aculeata excl. Formicidae from pitfall traps in the Calluna Heathlands of Northwest Germany. *ABH Naturwiss. Ver. Bremen.*, 41(2), 111-130.
- Roig-Alsina, A. (1991). Cladistic analysis of the Nomadinae s. str. with description of a new genus (Hymenoptera: Anthophoridae). *J. Kansas Entomol. Soc.*, 64(1), 23-37.
- Roig-Alsina, A. (1990). *Coelioxoides* Cresson, a parasitic genus of Tetrapediini (Hymenoptera: Apoidea). *J. Kansas Entomol. Soc.*, 63(2), 279-287.
- Roman'Kova, T. G. (1988). A new far eastern bee genus (*Lasanthidium*) of the tribe Anthidiini (Hymenoptera, Apoidea, Megachilidae). *Vestnik Zoologii*, (4), 25-30.
- Roseler, P. F. (1987). Grossenpolymorphismus, Geschlechtsregulation und Stabilisierung der Kasten im Hummelvolk. In: G. H. Schmidt, Sozialpolymorphismus bei Insekten. (pp. 298-335). Stuttgart: Wissenschaftliche Verlagsgesellschaft MBH.
- Roseler, P. F., & Honk, C. G. J. van. (1990). Castes and reproduction in bumblebees. In: W. Engels, Social Insects: an Evolutionary Approach to Castes and Reproduction. (pp. 147- 166). New York: Springer-Verlag.
- Roth, E. (1990). Beiträge der 3. Arbeitstagung zur Nutzung von Wildbestäubern in der Pflanzenzuchtung und Saatgutproduktion der DDR (Teil 2). 7. Erfahrungen mit der Haltung und dem Einsatz der roten Mauerbiene (*Osmia rufa*) in Kohlbefruchtsfrüppen. *Wiss Z. Martin Luther Univ. Halles-Wittenberg Math-Naturwiss Reihe*, 39(5), 11-14.
- Roubik, D. W. (1990). Mate location and mate competition in males of stingless bees (Hymenoptera: Apidae: Meliponinae). *Entomol. Gen.*, 15(2), 115-120.
- Roubik, D. W. (1990). A mixed colony of *Eulaema* (Hymenoptera: Apidae), natural enemies, and limits to sociality. *J. Kansas Entomol. Soc.*, 63(1), 150-157.
- Roubik, D. W. (1990). Niche preemption in tropical bee communities: a comparision of neotropical and Malesian faunas. In: S. F. Sakagami, R. I. Ohgushi, & D. W. Roubik, Natural History of Social Wasps and Bees in Equatorial Sumatra. (pp. 245-257). Sapporo, Japan: Hokkaido University Press.
- Rozen Jr., J. G. (1991). Nesting biology and mature larva of the bee *Idiomelissodes duplocincta* (Hymenoptera: Anthophoridae: Eucerini). *Am. Mus. Novit.*, (3012), 11pp.
- Rozen Jr., J. G. (1990). Pupa of the bee *Pararhophites orobinus* (Hymenoptera: Apoidea: Megachilidae). *J. N. Y. Entomol. Soc.*, 98(3), 379-382.
- Rozen Jr., J. G., & Buchmann, S. L. (1990). Nesting biology and immature stages of the bees *Centris caesalpiniae*, *Centris pallida* and the cleptoparasite *Ericrosis lata* (Hymenoptera: Apoidea: Anthophoridae). *Am. Mus. Novit.*, (2985), 30pp.
- Rozen Jr., J. G., & McGinley, R. J. (1991). Biology and larvae of the cleptoparasitic bee *Townsendiella pulchra* and nesting biology of host *Hesperapis larreae* (Hymenoptera: Apoidea). *Am. Mus. Novit.*, (3005), 11pp.

- Rozen Jr., J. G., & Roig-Alsina, A. (1991). Biology, larvae, and oocytes of the parasitic bee tribe Caenoprosopidini (Hymenoptera: Anthophoridae: Nomadinae). Am. Mus. Novit., (3004), 10pp.
- Rozen Jr., J. G., & Michener, C. D. (1988). Nests and immature stages of the bee *Paratetrapedia swainsonae* (Hymenoptera: Anthophoridae). Am. Mus. Nov., (2909), 1-13.
- Rust, R. W. (1990). Spatial and temporal heterogeneity of pollen foraging in *Osmia lignaria propinqua* (Hymenoptera: Megachilidae). Environ. Entomol., 19(2), 332-338.
- Ruszkowski, A., Bilinski, M., & Kaczmarcza, K. (1988). Food plants and economic importance of bees of the family Melittidae, and new species of *Melitta* (Apoidea, Melittidae). Pszczelnicze Zeszyty Naukowe, 32, 111-134.
- Ruszkowski, A., Bilinski, M., & Kaczmarcza, K. (1988). The presence of bumble bees in protected and unprotected environments in the landscape park of Kazimierz and its neighborhood. Pszczelnicze Zeszyty Naukowe, 32, 135-168.
- Ruz, L. (1991). Classification and phylogenetic relationships of the panurgine bees: the Calliopsini and allies (Hymenoptera: Andrenidae). Univ. Kansas Sci. Bull., 54(7), 210-256.
- Sakagami, S. F. (1991). The halictine bees of Sri Lanka and vicinity II. (*Nesohalictus*: Hymenoptera: Halictidae). Zool. Sci. (Tokyo), 8(1), 169-178.
- Sakagami, S. F. (1989). How does the myrmecophilous stingless bee *Trigona* (*Trigonella*) *moorei*, invade ants nests? Insectarium, 26(6), 164-175.
- Sakagami, S. F. (1990). *Lasioglossum nupricola*, a glacial age relic. Insectarium, 27(8), 260-269.
- Sakagami, S. F. (1987). Sozialstruktur und Polymorphismus bei Furchen- oder Schmalbienen (Halictidae). In: G. H. Schmidt, Sozialpolymorphismus bei Insekten. (pp. 257-293). Stuttgart: Wissenschaftliche Verlagsgesellschaft MBH.
- Sakagami, S. F., & Fukuda, H. (1989). Nest founding and nest survival in a eusocial halictine bee, *Lasioglossum duplex*: additional observations. Res. Popul. Ecol. (Kyoto), 31(1), 139-151.
- Sakagami, S. F., & Inoue, T. (1990). Oviposition behavior of two Sumatran stingless bees, *Trigona (Tetragonula) laeviceps* and *T. (T.) fuscobalteata*. In: S. F. Sakagami, R. I. Ohgushi, & D. W. Roubik, Natural History of Social Wasps and Bees in Equatorial Sumatra. (pp. 201-217). Sapporo, Japan: Hokkaido University Press.
- Sakagami, S. F., Inoue, T., & Salmah, S. (1990). Stingless bees of Central Sumatra. In: S. F. Sakagami, R. I. Ohgushi, & D. W. Roubik, Natural History of Social Wasps and Bees in Equatorial Sumatra. (pp. 125-137). Sapporo, Japan: Hokkaido University Press.
- Sakagami, S. F., & Maeta, Y. (1989). Compatibility and incompatibility of solitary life with eusociality in two normally solitary bees *Ceratina japonica* and *Ceratina okinawana* (Hymenoptera, Apoidea), with notes on the incipient phase of eusociality. Jpn. J. Entomol., 57(2), 417-439.
- Sakagami, S. F., & Maeta, Y. (1990). *Lasioglossum primavera* sp. nov., a Japanese halictine bee which overwinters in both female and male adults (Hymenoptera, Halictidae). Bull. Fac. Agric. Shimane Univ., (24), 52-59.
- Sakagami, S. F., & Maeta, Y. (1990). *Trigona (Tetragonisca) angustula*, a distinguished stingless bee species. Insectarium, 27(1), 13-14.
- Sakagami, S. F., & Munakata, M. (1990). *Lasioglossum blakistoni* sp. nov., the northernmost representative of the paleotropic subgenus *Ctenonomia* (Insecta, Hymenoptera, Halictidae). Zool. Sci. (Tokyo), 7(5), 985-988.
- Sakagami, S. F., Kato, M., & Itino, T. (1991). *Trinchostoma (Diagonozus) asanum* sp. nov.: halictine bees from Sumatra, with some observations on its oligotrophy to *Impatiens*. Topics, 1, 49-58.
- Salmah, S., Inoue, T., & Sakagami, S. F. (1990). An analysis of apid bee richness (Apidae) in central Sumatra. In: S. F. Sakagami, R. I. Ohgushi, & D. W. Roubik, Natural History of Social Wasps and Bees in Equatorial Sumatra. (pp. 139-174). Sapporo, Japan: Hokkaido University Press.
- Sazima, M., & Sazima, M. (1989). Flower visits by carpenter bees and stingless bees, their interactions and consequences for the pollination of yellow passionflower. Revista Brasileira de Entomologia, 33(1), 109-118.

- Sazima, M., & Sazima, I. (1989). Oil-gathering bees visit flowers of eglandular morphs of oil-producing Malpighiaceae. *Botanica Acta*, 102, 106-111.
- Schemske, D. W., & Horvitz, C. C. (1989). Temporal variation in selection on a floral character. *Evolution*, 43(2), 461-465.
- Schmid-Hempel, P., Muller, C., Schmid-Hempel, R., & Shykoff, J. A. (1990). Frequency and ecological correlates of parasitism by conopid flies (Conopidae, Diptera) in populations of bumblebees. *Insectes Soc.*, 37(1), 14-30.
- Schmid-Hempel, P., & Schmid-Hempel, R. (1990). Endoparasitic larvae of conopid flies alter pollination behavior of bumblebees. *Naturwissenschaften*, 77(9), 450-452.
- Schmid-Hempel, R., & Muller, C. B. (1991). Do parasitized bumblebees forage for their colony? *Animal behaviour*, 41(5), 910-912.
- Schmid-Hempel, R., & Schmid-Hempel, P. (1991). Endoparasitic flies pollen-collected by bumblebees and a potential host-parasite conflict. *Oecologia (Heidelb.)*, 87(2), 227-232.
- Schmitt, U. (1990). Hydrocarbons in tarsal glands of *Bombus terrestris*. *Experientia*, 46(10), 1080-1082.
- Schneider, S. S. (1991). Modulation of queen activity by the vibration dance in swarming colonies of the African honey bee *Apis mellifera scutellata* (Hymenoptera: Apidae). *Insectes Sociaux*, 38(2), 269-279.
- Scholl, A., Obrecht, E., & Owen, R. E. (1990). The genetic relationship between *Bombus moderatus* Cresson and the *Bombus lucorum* Auct. species complex (Hymenoptera: Apidae). *Can. J. Zool.*, 68 (11), 2264-2268.
- Schonitzer, K., & Seifert, P. (1990). Anatomy and ultrastructure of the salvary gland in the thorax of the honeybee worker, *Apis mellifera* (Insecta, Hymenoptera). *Zoomorphology*, 109, 211-222.
- Schonitzer, K., & Klinksik, C. (1990). The ethology of the solitary bee *Andrena nycthemera* Imhoff, 1866 (Hymenoptera, Apoidea). *Entomofauna Zeitschrift fur Entomologie*, 23(1), 377-428.
- Schonitzer, K., & Schmid, S. (1990). Antennales Sensillenmuster bei Verschiedenen arten von *Andrena* (Hymenoptera, Andrenidae). *Mitt. Dtsch. Ges. Allg. Angew. Ent.*, 7, 475-478.
- Schonitzer, v. K., & Klinksik, C. (1990). Individuell unterschiedlicher Lebenslauf bei der Sandbiene *Andrena nycthemera* Imhoff (Hymenoptera, Apoidea). *Nachr. Bl. Bayer. Ent.*, 39(4), 116-121.
- Shelly, T. E., Buchmann, S. L., Villalobos, E. M., & O'Rourke, M. K. (1991). Colony ergonomics for a desert-dwelling bumblebee species (Hymenoptera: Apidae). *Ecol. Entomol.*, 16 (3), 361-370.
- Shykoff, J. A., & Schmid-Hempel, P. (1991). Genetic relatedness and eusociality: parasite-mediated selection on the genetic composition of groups. *Behav. Ecol. Sociobiol.*, 28(5), 371-376.
- Shykoff, J. A., & Schmid-Hempel, P. (1991). Incidence and effects of four parasites in natural populations in bumble bees in Switzerland. *Apidologie*, 22(2), 117-126.
- Shykoff, J. A., & Schmid-Hempel, P. (1991). Parasites and the advantage of genetic variability within social insect colonies. *Proc. R. Soc. Lond. Ser. Biol. Sci.*, 243(1306), 55-58.
- Silva, M. F. da, Miranda, I. P. de A., & Barbosa, E. M. (1986/ 1987). Aspects of the pollination of the african oil palm (*Elaeis guineensis*) and the american oil palm (*Elaeis oleifera*). *Acta Amazonica*, 16-17, 209-218.
- Singh, A., & Tiwari, V. K. (1988). Caste differentiation in *Halictus ducalis* Bingham (Hymenoptera: Halictidae). *J. Advanced Zool.*, 9(1), 74-75.
- Skutch, A. F. (1989). Flowering and seed production of *Fischeria funebris* Asclepiadaceae. *Brenesia*, (30), 13-18.
- Smith, B. H., & Weller, C. (1989). Social competition among gynes in halictine bees: the influence of bee size and pheromones on behavior. *J. Insect Behav.*, 2(3), 397-411.
- Smith, D. R. (1991). African bees in the Americas: insights from biogeography and genetics. *Trends Ecol. Evol.*, 6(1), 17-21.
- Snelling, R. R. (1990). A review of the native North American bees of the genus *Chalicodoma* (Hymenoptera: Megachilidae). *Contrib. Sci.*, (421), 1-39.
- Sommeijer, M. J., & de Bruijn, L. L. M. (1988). *Melipona trinitatis* and *Melipona favosa*, the only two species of the genus *Melipona* in Trinidad. *Uitg. Natuurwet. Studiekring Suriname Ned Antillen*, (123), 75-82.

- Spangler, H. G., & Buchmann, S. L. (1991). Effects of temperature on wingbeat frequency in the solitary bee *Centris caesalpiniae* (Anthophoridae: Hymenoptera). J. Kansas Entomol. Soc., 64(1), 107-109.
- Spivak, M., Fletcher, D. J. C., & Breed, M. D. (Eds.) (1991). The "African" Honey Bee. Boulder: Westview Press, 435 pp.
- Stark, R. E. (1989). Beobachtungen zur Nestgründung und Brutbiologie der Holzbiene *Xylocopa sulcatipes* Maa (Apoidea: Anthophoridae). Mitt. Dtsch. Ges. Allg. Angew. Entomol., 7(1), 252-256.
- Stark, R. E., Hefetz, A., Gerling, D., & Velthius, H. H. W. (1990). Reproductive competition involving oophagy in the socially nesting bee *Xylocopa sulcatipes*. Naturwissenschaften, 77(1), 38-40.
- Starr, C. K., & Geronimo, J. G. (1990). Habitat and foraging observations on an oriental bumble bee (Hymenoptera: Apidae). Ethol. Ecol. Evol., 2(4), 373-379.
- Steiner, K. E., & Whitehead, V. B. (1990). Leg length evolution among oil-collecting bees. In: Proceedings of the Fourth International Congress of Systematic and Evolutionary Biology; College Park, Maryland, USA, July 1-7, 1990.
- Steinmann, E. (1990). Zur Nahorientierung der solitären Sandbiene *Andrena vaga* Panzer 1799 (Hymenoptera, Apoidea) am Nesteingang. Mitt. Schweiz. Entomol. Ges., 63(1-2), 77-80.
- Steinmann, E., & Menzel, R. (1990). Lernversuche mit der Einsiedlerbiene *Osmia rufa* (Linnaeus, 1758) (Hymenoptera, Apoidea). Mitt. Schweiz. Entomol. Ges., 63, 99-103.
- Stephen, W. P., & Fichter, B. L. (1990). Chalkbrood (*Ascophaea aggregata*) resistance in the leafcutting bee (*Megachile rotundata*). 2. Random matings of resistant lines to wild type. Apidologie, 21(3), 221-231.
- Stephen, W. P., & Fichter, B. L. (1990). Chalkbrood *Ascophaea aggregata* resistance in the leafcutting bee *Megachile rotundata* I. Challenge of selected lines. Apidologie, 21 (3), 209-220.
- Stern, D. L., & Dudley, R. (1991). Wing buzzing by male orchid bees, *Eulaema meriana* (Hymenoptera: Apidae). J. Kansas Entomol. Soc., 64(1), 88-94.
- Sugiura, N., & Maeta, Y. (1989). Parental investment and offspring sex ratio in a solitary mason bee, *Osmia cornifrons* (Radoszkowski) (Hymenoptera, Megachilidae). Jpn. J. Entomol., 57(4), 861-875.
- Surholt, B., Grieve, H., Baal, T., & Bertsch, A. (1990). Non-shivering thermogenesis in asynchronous flight muscles of bumblebees. Comparative studies on males of *Bombus terrestris*, *Xylocopa sulcatipes* and *Acherontia atropos*. Comp. Biochem. Physiol. A Comp. Physiol., 97(4), 493-499.
- Surholt, B., Grieve, H., Baal, T., & Bertsch, A. (1991). Warm-up and substrate cycling in flight muscles of male bumblebees, *Bombus terrestris*. Comp. Biochem. Physiol. A Comp. Physiol., 98(2), 299-303.
- Sutcliffe, G. H., & Plowright, R. C. (1990). The effects of pollen availability on development time in the bumble bee *Bombus terricola* K. (Hymenoptera: Apoidea). Can J. Zool., 68(6), 1120-1123.
- Szabo, T. I., & Lefkovitch, L. P. (1991). Development of overwintered honey bee colonies with one- and two-year-old queens. Bee Science, 1(3), 144-150.
- Tealdi, C. (1989). Bees (*Apis dorsata*) in Malaysia. Ape Nostra Amica, 11(4), 15-19.
- Tengo, J., Agren, L., Baur, B., Isaksson, R., Liljefors, T., Mori, K., Konig, W., & Franke, W. (1990). *Andrena wilkella* male bees discriminate between enantiomers of cephalic secretion components. J. Chem. Ecol., 16(2), 429-441.
- Teras, I. (1990). Odors in the life of bumblebees. Luonnon Tutkija, 94(5), 204-211.
- Thalmann, U., & Dorn, M. (1990). Beiträge der 3. Arbeitstagung zur Nutzung von Wildbestäubern in der Pflanzenzuchtung und Saatgutproduktion der DDR (Teil 2). 8. Die Haltung der Pelzbiene, *Anthophora acervorum* (L.), und ihr Einsatz zur Nutzpflanzenbestäubung. Wiss Z. Martin Luther Univ. Halle-Wittenberg Math-Naturwiss Reihe, 39(5), 15-21.
- Togashi, I. (1990). Hymenopterous insects settling in cottage with a thatched roof in Shiramine-Mura Ishikawa prefecture Japan part 1. Bull. Biogeogr. Soc. Jpn., 45(1-22), 111-116.
- Torchio, P. F. (1990). *Osmia ribifloris*, a native bee species developed as a commercially managed pollinator of highbush blueberry (Hymenoptera: Megachilidae). J. Kansas Entomol. Soc., 63(3), 427-436.
- Torchio, P. F. (1991). Use of *Osmia lignaria propinqua* (Hymenoptera: Megachilidae) as a mobile pollinator of orchard crops. Environ. Entomol., 20(2), 590-596.
- Toro, H., Chiappa, E., Ruz, L., & Cabezas, V. (1991). Comportamiento reproductivo de *Centris mixta* Tamarugalensis (Hymenoptera, Anthophoridae). 1 parte. Acta Entomol. Chil., 16, 97-112.

- Toro, H., & Fritz, M. (1991). Contribucion al conocimiento de *Dasycoelioxys* Mitchell (Hymenoptera, Apoidea, Megachilidae). *Acta Entomol. Chil.*, 16, 69-80.
- Torres, F., Gayubo, S. F., & Asensio, E. (1989). Efecto de la presion urbana sobre abejas y avispas (Hymenoptera, Aculeata) en Salamanca. V. Superfamilia Apoidea. *Comun. Inst. Nac. Invest. Agro. Ser. Recur. Nat.*, 53, 1-49.
- Urban, D. (1989). Duas especies novas do genero *Trichocerapis* Cockerell, 1904 (Hymenoptera, Apoidea). *Rev. Bras. Zool.*, 6(3), 457-462.
- Urban, D. (1988). New species of *Melissoptila* from Cerrado of Parana Brazil (Hymenoptera: Apoidea). *Acta Biol. Parana.*, 17(1-4), 1-10.
- Urbani, C. B. (1989). On a singular case of kin selection theory among Hymenoptera justified fratricide and the natural way to avoid it. *Ethol. Ecol. Evol.*, 1(4), 373-378.
- Van Asperen de Boer, J. R. J. (1990). *Bombus krusemani* - a new bumblebee species from Guatemala (Hymenoptera: Apidae). *Ent. Ber., Amst.*, 50(1), 1-3.
- Van der Zanden, G. (1989). New or little known species and subspecies of paleartic megachilid species (Insecta: Hymenoptera: Apoidea: Megachilinae). *Entomol. Abh. (Dres.)*, 53(1-7), 71-86.
- Van der Zanden, G. (1990). Nomenclature changes for some paleartic species of the family Megachilidae (Insecta: Hymenoptera: Apoidea). *Reichenbachia*, 28(1-19), 51-54.
- Van der Blom, J. (1991). Social regulation of egg-laying by queenless honeybee workers (*Apis mellifera* L.). *Behav. Ecol. Sociobiol.*, 29(5), 341-346.
- Van Doorn, A., & Chrambach, A. (1989). Retinue behavior in bumblebee workers (*Bombus terrestris* L.). *J. Apic. Res.*, 28(2), 66-70.
- Vardy, C. R. (1989). The biology of the bee *Chelostoma florisomne* (Linnaeus) in Britain (Hymenoptera: Megachilinae). *Entomologist*, 108(3), 167-175.
- Velthuis, H. H. W. (1988). Between hollow and honeycomb. *Endeavour*, 12(1), 2-7.
- Velthuis, H. H. W. (1989). Chemical signals and dominance communication in the honeybee *Apis mellifera* (Hymenoptera: Apidae). *Entomol. Gener.*, 15(2), 83-90.
- Velthuis, H. H. W. (1989). Regulation of honeybee reproduction at colony level. *Honeybee Science*, 10(4), 155-159.
- Velthuis, H. H. W., & Cobb, L. (1991). Pollination of *Primula* in a greenhouse using bumblebees. *Acta Horticulture* 288, 6th Pollination Symposium, p.199-203.
- Velthuis, H. H. W., Ruttner, F., & Crewe, R. M. (1990). Differentiation in reproductive physiology and behavior during the development of laying worker honey bees. In: W. Engels, *Social Insects: An Evolutionary Approach to Castes and Reproduction*. (pp. 231-244). Berlin: Springer-Verlag.
- Velthuis, H. H. W., & Van der Kerk, A. (1988). Age, environment, and genes in relation to the mandibular gland secretion of pure and hybrid *Apis mellifera capensis* worker bees. In: G. R. Needham, *Africanised Honey Bees and Bee Mites*. (pp. 80-86). Cichester, England: Ellis Horwood Ltd.
- Villa, J. D., & Weiss, M. R. (1990). Observations on the use of visual and olfactory cues by *Trigona* spp. foragers. *Apidologie*, 21(6), 541-545.
- Villalobos, E. M., & Shelly, T. E. (1991). Correlates of male mating success in two species of *Anthidium* bees (Hymenoptera: Megachilidae). *Behav. Ecol. Sociobiol.*, 29 (1), 47-54.
- Vinson, S. B., & Frankie, G. W. (1990). Territorial and mating behavior of *Xylocopa fimbriata* F. and *Xylocopa gualanensis* Cockerell from Costa Rica. *J. Insect Behav.*, 3(1), 13-32.
- Vinson, S. B., & Frankie, G. W. (1991). The oil baron bees of Lomas Barbudal. *Bee Line News and Bulletin; Friends of Lomas Barbudal*, 5(1,2,&3), 16-17.
- Vogel, S. (1987). History of the Malpighiaceae in the light of pollination ecology. *Mem. N.Y. Bot. Gard.*, 55, 130-142.
- Waddington, K. D. (1988). Body size, individual behavior and social behavior in honey bees. R. L. Jeanne, *Interindividual Behavioral Variability in Social Insects*. (pp. 385-418). Boulder & London: Westview Press.
- Walch, H., & Straub, M. (1990). Arthropod fauna in farms with and without the use of insecticides against the second generation of the grape fruit and berry moth. In: W. Laux, *Mitteilungen aus der Biologischen Bundesanstalt fur Land-und Forstwirtschaft Berlin-Dahlem*, Heft 266. 47. Berlin, Germany: Deutsche Pflanzenschutz-Tagung.

- Walsh, S. L. (1990). Collection of *Dufourea versatilis* new record (Hymenoptera: Halictidae) from Idaho USA. *Entomol. News*, 101(1), 67-68.
- Warncke, K. (1989). Habitats, dates and labelling of bees collected (in Turkestan) by A. Fedtschenko in 1868-71. *Linzer Biologische Beitrage*, 21(1), 3-14.
- Waser, N. M., & Price, M. V. (1990). Pollination efficiency and effectiveness of bumble bees and hummingbirds visiting *Delphinium nelsonii*. *Collect. Bot.*, 19, 9-20.
- Watson, R. H., & Van Ark, H. (1989). The effect of temperature on the developmental rate of the immature stages of large carpenter bees *Xylocopa* spp. (Hymenoptera: Anthophoridae). *J. Entomol. Soc. South Afr.*, 52(1), 119-128.
- Wcislo, W. T. (1990). A new species of *Lasioglossum* from Costa Rica (Hymenoptera: Halictidae). *J. Kansas Entomol. Soc.*, 63(3), 450-453.
- We, Y. R. (1990). A study on Chinese *Dufourea* with descriptions of five new species (Hymenoptera: Apoidea: Halictidae). *Acta Entomol. Sin.*, 33(4), 466-475.
- Wehner, R. (1990). Do insects have cognitive maps? *Annu. Rev. Neurosci.*, 13, 403-414.
- Wehner, R. (1989). The Hymenopteran skylight compass: matched filtering and parallel coding. *J. Experimental Biol.*, 146, 63-85.
- Westerkamp, C. (1991). Honeybees are poor pollinators - why? *Pl. Syst. Evol.*, 177, 71-75.
- Widen, B., & Widen, M. (1990). Pollen limitation and distance-dependent fecundity in females of the clonal gynodioecious herb *Glechoma hederacea* (Lamiaceae). *Oecologia*, 83(2), 191-196.
- Williams, P. H. (1989). Bumble bees and their decline in Britain. Ilford, Great Britain: The Central Association of Bee-Keepers, pp.1-15.
- Williams, P. H. (1991). The bumble bees of the Kashmir Himalaya (Hymenoptera: Apidae, Bombini). *Bull. Br. Mus. nat. Hist. (Ent.)*, 60(1), 1-204.
- Wilson, E. O. (1990). Excellence in Ecology, 2. Success and Dominance in Ecosystems; the Case of the Social Insects. Oldendorf/Luhe, Germany: Ecology Institute.
- Wilson, H. D. (40). Gene flow in squash species. *Biosci. Am. Inst. Biol. Sci.*, 6(449-455).
- Winn, B. (1988). Importation and release of leafcutter bees (*Megachile rotundata*) in South Australia. In: J. W. Rhodes, *Bee Keeping in the Year 2000*. (pp. 209-210). Queensland, Australia: International Colour Productions Pty.
- Winston, M. L. (1991). The inside story internal colony dynamics of Africanized bees. In: M. Spivak, D. J. C. Fletcher, & M. D. Breed, *The "African" Honey Bee*. (pp. 201-212). Boulder, Colorado, USA; Oxford, England, UK: Westview Press.
- Wittmann, D., Radtke, R., Cure, J. R., & Schifino-Wittmann, M. T. (1990). Coevolved reproductive strategies in the oligoleptic bee *Callonychium petuniae* (Apoidea: Andrenidae) and three purple flowered *Petunia* spp. Solanaceae in southern Brazil. *Z. Zool. Syst. Evolutionsforsch*, 28(3), 157-165.
- Wittmann, D., Radtke, R., Zeil, J., Lubke, G., & Franke, W. (1990). Robber bees (*Lestrimelitta limao*) and their host: chemical and visual cues in nest defense by *Trigona (Tetragonisca) angustula* (Apidae: Meliponinae). *J. Chem. Ecol.*, 16(2), 631-641.
- Wittmann, D., Radtke, R., Hoffmann, M., & Blochtein, B. (1989). Seasonality and seasonal changes in preferences for scent baits in *Euplusia violacea* in Rio Grande do Sul/Brazil (Hymenoptera: Apidae: Euglossini). *Entomol. Gen.*, 14(3/4), 217-221.
- Wolfe, L. M., & Barrett, S. C. H. (1989). Patterns of pollen removal and deposition in tristylous *Pontederia cordata* L. (Pontederiaceae). *Biol. J. Linnean Soc.*, 36, 317-329.
- Wongsiri, S., Lekprayoon, C., & Pothichot, S. (1989). Efficiency of crossing Chinese strain *Apis cerana cerana* and Thai strain *Apis indica* (Apidae: Hymenoptera) by artificial insemination. *Honeybee Sci.*, 10(3), 112-114.
- Wongsiri, S., Limbipichai, K., Tangkanasing, P., Marden, M., Rinderer, T., Sylvester, H. A., Koeniger, G., & Otis, G. (1990). Evidence of reproductive isolation confirms that *Apis andreniformis* (Smith, 1858) is a separate species from sympatric *Apis florea* (Fabricius, 1787). *Apidologie*, 21(1), 47-52.
- Woyke, J., & Jasinski, Z. (1990). Effect of the number of attendant worker bees on the initiation of egg laying by instrumentally inseminated queens kept in small nuclei. *J. Apic. Res.*, 29(2), 101-106.

- Wu, Y. R. (1988). Four new species of bees from China (Hymenoptera, Apoidea) (*Melitta* spp.; *Anthomegilla* spp.). *Acta Zootaxonomica Sinica*, 13(1), 67-71.
- Wu, Y. R. (1988). A new species of *Anthophora* (*A. antennalis*) from China (Hymenoptera: Apoidea, Anthophoridae). *Acta Entomologica Sinica*, 31(2), 210-212.
- Wu, Y. R. (1990). Descriptions of nine new species of Apoidea from inner Mongolia China. *Entomotaxonomia*, 12(3-4), 243-252.
- Wu, Y. R. (1990). A study on Chinese *Dufourea* with descriptions of five new species (Hymenoptera: Apoidea: Halictidae). *Acta Entomol. Sin.*, 33(4), 466-475.
- Wuest, J. (1990). Les 'outils' de l'abeille. *Mus. Geneve*, (307), 8-13.
- Xiong, L., & Xing, Z. (1988). Pollinating bees for *Amomum villosum* in Yunnan. *Chin. J. Trop. Crops*, 8(2), 75-81.
- Yanega, D. (1990). Philopatry and nest founding in a primitively social bee *Halictus rubicundus*. *Behav. Ecol. Sociobiol.*, 27(1), 37-42.
- Yeskov, Y. K. (1990). Mechanisms of perception by honey bee of low-frequency electric fields. *Zoological Journal*, 69(5), 53-59.
- Yeskov, E. K., & Mironov, G. A. (1990). Low-frequency electric field and concomitant physical factors influence upon bees, comparative analysis. *Ecologiya* (Sverdlovsk), (6), 81-84.
- Yong, H. S. (1991). Biochemical genetic differentiation in the *Trigona atripes* species group of Malaysian stingless bees (Insecta: Hymenoptera: Apidae). *Comp. Biochem. Physiol.*, 99B(3), 625-628.
- Young, C. G., & Owen, R. E. (1989). Foraging activity of bumble bee, *Bombus* spp., workers on yellow *Hedysarum*, *Hedysarum sulphurescens*, in a subalpine meadow. *Can. Field Nat.*, 103(3), 341-347.
- Zakhvatkin, V. K. (1990). Species composition and features of the formation of complexes of bees - pollinators of lucerne in the Alma Ata Region. *Tr. Inst. Zool. Akad. Nauk. Kaz. SSR*, 45, 159-163.
- Zhang, J. F. (1990). New fossil species of Apoidea Insecta Hymenoptera. *Acta Zootaxonomica*, 15(1), 83-91.
- Zhilyaev, G. G. (1989). Formation of population borders of entomophilous plants in connection with pollination. *Zh. Obshch. Biol.*, 50(5), 646-654.
- Zhou, Y. F., Luo, Y. X., Chen, H. S., & Lai, Y. S. (1989). Occurrence and harmfulness of *Galleria mellonella* L. (Lep.: Pyralidae). *Natural Enemies of Insects*, 11(2), 87-93.

ANNOUNCING
**JOURNAL OF
HYMENOPTERA RESEARCH**

Published by
The International Society of Hymenopterists

A new journal devoted to all aspects of research on the Hymenoptera
Subject matter may include biology, biocontrol, behavior, ecology,
systematics, taxonomy, genetics and morphology

Anticipated contents of first volume to be issued in early 1992:

- B. Alexander - An exploratory analysis of cladistic relationships within the superfamily Apoidea, with special reference to sphecid wasps
R. Bohart - A synopsis of the genus *Oxybelus* in Middle America
J. Carpenter & D. Brothers - Aculeate phylogeny
F. Gess & S. Gess - Nesting ethology of *Celonites* and *Quartinia*
F. Gess - A new species of *Celonites*
E. Grissell - A revision of the genus *Perissocentrus* (Torymidae)
L. Kimsey - Functional morphology of the abdomen and phylogeny of chrysidid wasps
B. Norden, K. Krombein & J. Steinberg - Mating behavior, male allometry and nesting biology of *Perdita graenicheri*
W. Pulawski & H. Court - Revision of the genus *Hingstoniella*
D. Rosen & J. Woolley - Biological and taxonomic studies of *Chartocerus subaeneus* (Signiphoridae)
D. Stoltz & J. Whittfield - Polydnaviruses and parasitoids: an introduction for Hymenopterists
M. Sharkey & D. Wahl - Cladistic analysis of ichneumonoid families
A. Menke - Mole cricket hunters of the genus *Larva* in the New World
F. Kurczewski, M. O'Brien & M. Spofford - Nesting behavior of *Podalonia robusta* (Sphecidae)
J. Aeschlimann & R. Hughes - Collecting *Aphelinus* spp. in southwestern USSR for pre-emptive biological control of Russian wheat aphid in Australia
D. Queller, J. Strassmann & C. Hughes - Genetic relatedness and population structure in primitively eusocial wasps in the genus *Mischocyttarus*
F. Jachmann & J van den Assam - Selective perception in courting male parasitic wasps
G. Gibson - Evolutionary relationships and classification of the Symphyta
J. Whittfield - Phylogeny of non-Aculeate Hymenoptera
A. Austin, R. Wharton & S. Cameron - Molecular techniques in Hymenoptera systematics
D. Brothers - The first Mesozoic Vespidae from the Southern Hemisphere, Botswana

For information and instructions to authors, write to

David R. Smith, Editor
Systematic Entomology Laboratory
U. S. Department of Agriculture
c/o National Museum of Natural History NHB-168
Washington, DC 20560

Subscriptions for libraries are \$50/year. Send order to the editor (see address above).

Publication in the *Journal of Hymenoptera Research* is FREE to members of The International Society of Hymenopterists. To join ISH, complete the membership application below.

Application for membership in The International Society of Hymenopterists

Name _____ Phone _____
Address _____

Area of interest _____

Membership includes a subscription to and free publication privileges in the *Journal of Hymenoptera Research*. Payment may be made in US or Canadian dollars.

Categories of membership (check one):

Regular ____ \$25 US (\$29 Can) Student ____ \$15 US (\$18 Can) Sustaining ____ \$50 US (\$58 Can)

Also enclosed is my tax deductible contribution of \$ _____ to help support the operation of ISH

Send application with payment to:

Gary Gibson, Treasurer
Biological Resources Division/CLBRR
Agriculture Canada
K. W. Neatby Building
Ottawa, Ontario, Canada K1A 0C6