

A FORUM FOR ACULEATE WASP RESEARCHERS

# \$\$\$ SEND MONEY \$\$\$ PLEASE!

As of December 16 | have not received my budget for FY94, but the feeling of leadership is that in all likelihood it will be very small, and thus insufficient to pay for reproduction of one issue of Sphecos. Thus your editor is going to pay for this issue out of personal funds (Ouch!). In order to recoup my outlay (591 dollars) for Sphecos 26, I would appreciate contributions from the readership. If each of you sends \$5 (or whatever you can afford) that should defray my expense for issue 26, and hopefully provide money for issue 27, and possibly also 28. I know of no easier way to resolve this problem, and I hope that most of you regard this rag as sufficiently wonderful that you will come through with monetary aid. If you send your contribution with a personal check, please make it payable to Arnold Menke. Thank you!

# NOTICE

We now have a link to the Internet. Terry has a logon with the USDA computer at Beltesville, Maryland, and can receive E-MAIL and download files to his own computer. Those of you who wish to contribute to **Sphecos** with an article written on a computer and have access to the Internet can choose to send it via E-MAIL instead of mailing us a floppy disk. We can handle both ASCII files and word processing documents, just as long as we know the format of your document. Diacritics sometimes don't convert very well when going from a ARNOLD S. MENKE, Editor TERRY NUHN, Assistant Editor Systematic Entomology Laboratory Agricultural Research Service, USDA c/o National Museum of Natural History Smithsonian Institution, Washington, DC 20560 FAX: (202) 786-9422 Phone: (202) 382-1803 E-MAIL (Internet): TNUHN@asrr.arsusda.gov

DOS machine to a Macintosh, so you might want to include an explanatory note to help us with these. Terry's address is:

TNUHN @ASRR.ASRUSDA.GOV.

# **RESEARCH NEWS**

Alexander Antropov (Zoological Museum of Moscow Lomonosov State Univ., Herzon Street, Mowcow, 103009, Russia) reports: "I intend to submit an application for a long-term grant from the International Science Foundation in September [1993]. I am planning to review the Palearctic Belomicrus [one paper is listed in Recent Literature and two others are in press] as a project. The ISF conditions imply that the work is to be completed in two years and mainly in the territory of the former Soviet Union, and I want to use this opportunity to visit some places in the Central Asian republics as well as the BM(NH) to study type material." [Alexander is also applying to the Smithsonian Institution for a short term visitor grant so that he can study North American *Trypoxylon* (*Trypoxylon*) as a part of his revision of the Holarctic species - editor.]

**Celso Oliveira Azevedo** (Universidad Federal do Espirito Santo, Dept. Biologia, Av. Marechal Campos 1468, Maruipe, 29040-090 Vitória, ES, Brasil) writes: "I am now working as a teacher of zoology at the Universidad Federal do Espirito Santo and am presently working on my Ph.D. program, a systematic and biogeographical revision of the neotropical *Dissomphalus* (Hymenoptera, Bethylidae)."

Eduardas Budrys (Institute of Ecology, Akademijos 2, Vilnius 2600, Lithuanian Republic) is developing a computer database of the Pemphredoninae: "My database already takes up over 5 MB of disk space (I say "my" but most of the data, especially bibliography are really Wojciech's [Pulawski]). It consists of 19 relationally linked data tables (SPECIES, NOMSPEC, GENERA, NOMGEN, AUTHORS, MUSEUMS. SPECREF, GENREF, CHARS, ITEMS, MATERIAL, etc.) and contains not only the bibliography for each species of Pemphredoninae, but also addresses of authors, data about museums and collections, types, collected material, collecting localities, morphometric data of measured specimens, list of characters and states of each character, and even descriptions of species in the DELTA program format (the last can be used for cladistic analysis, too). The database is in PARADOX 4.0 format. Morphometric data are analysed using the spreadsheet QUATTRO PRO for WINDOWS (the last program can immediately input data from PARADOX data tables and output them back). I hope that by joint efforts of sphecidists a data base of all Sphecidae can be created. I would like to contact everybody working on wasp taxonomic databases to standardize data structure, etc. [You might be interested in the *BIOTA* information system described on p. 4 - editor.]

**Derek Daly** (133 Linner Road, Speke, Liverpool L24 3OQ, Great Britain) is interested in *Vespula vulgaris*, particulary in foraging behaviour and relatedness among nestmates.

Julio Antonio Genaro (25 no. 466, Vedado 12300, Cuidad de La Habana, Cuba) writes: "I am describing a new species of *Lophostigma* (Mutillidae). This genus is known only from Central and South America. Also, I have other interesting mutillids that are perhaps new for the West Indies."

Edilberto Giannotti (Dept. Zoologia, Inst. Biociências, Univ. Estadual Paulista, Caixa postal 199, Rio Claro, SP, Brasil, 13.560-900) reports: "I am engaged in research on social wasps, mainly *Polistes* and *Mischocyttarus*, concerning their biological cycles, behavior, foraging, and their role in the ecosystem. I just defended my doctoral dissertation on *Polistes Ianio* in November, 1992."

Jean Leclercq (Faculté d'Agronomie de Gembloux, rue de Bois-de-Breux, 190, B 4020, Liege - Jupille, Belgium) says: "Last year and this year up to February, I was busy (too busy for anything else) with my book with Jacques Bitsch, Faune de France, Sphecidae I: Généralités, Crabroninae, and with two parallel papers recording data on the distribution of European Crabronini. The book has more than 300 pages and is due to be published in September, 1993, while the other two papers are due to appear before the end of 1993.

"Having written these papers, I spent much of my time last spring studying Crabronini, including hundreds of North American specimens, received on loan from the Snow Museum (Lawrence, Kansas) and from the Florida Deptartment of Agriculture (Gainesville); I got as far as getting quite a good idea of the N. American *Crossocerus*, based on an unpublished key Richard Miller prepared years ago and on information we exchanged when that unhappy colleague was so remarkably active. So American *Crossocerus* are also in my mind, but not for the coming months. I must also please Krombein with some Sri Lanka material."

Alex Rasnitsyn (Arthropoda Laboratory, Paleontological Institute, Russian Academy of Sciences, Profsoyuznaya Str. 123, Moscow, Russia, 117647) is still completely engaged in administering the program by Mr. George Soros to support biologists in the former Soviet Union, using 4,000 individual grants in biodiversity of \$500 each, to be distributed in open competition. This takes almost all his (Alex) time leaving practically none for research. However, Alex hopes that he will be able to finish this program by the New Year and to return to his earlier projects now in suspension. Among these are: (i) The History of Insects (two multiauthored volumes in English, with the first one reviewing the systematic, phylogenetic and paleontological history of the insect orders being nearly ready). (ii) Revision of the Cretaceous Sphecidae known from impression fossils, with numerous new descriptions (with Woj Pulawski, nearly half-ready). (iii) An analysis of the insect fauna of the Dominican amber based in particular on the Smithsonian collection (with Don Davis and possibly other contributors, nearly ready). Possibly (depending on the availability of a grant applied for to the International Science Foundation) he also has a new project: The Worldwide Database of the non-Quarternary Fossil Insects, to be completed by the staff of the Arthropoda Laboratory, Paleontological Institute, Russian Academy of Sciences, Moscow, Russia.

**Katsuji Tsuneki** (Asahigaoka 4-15, Mishima, Japan 411), who is now 85 years old, writes: "At present I am tackling a revision of *Sphecodes* in Japan (possibly my final work.)"

**Colin Vardy** (Dept. of Entomology, The Natural History Museum, Cromwell Road, London SW7 5BD UK) writes: "I would like to place on record my intention of revising the sphecid genus *Podium*. Readers will already be aware that **Servio Amarante** (São Paulo) is revising the closely related *Penepodium*. We will be comparing their biogeography etc. He wishes also to compare the phylogenetics of these two genera, together with *Dynatus* and *Trigonopsis*, since they form a natural group (formerly called 'Podiini') within the Sceliphrini. **Michael Ohl** (Kiel) wishes to take this further and work out the phylogenetics of the entire subfamily Sphecinae."

Rolf Witt (AG Terrestrische Okologie, Fachbereich 7, University Oldenburg, Postfach 2503,26129 Oldenburg, Germany) says: "In the next two years I'll be busy with my doctoral thesis, that deals with the distribution and indicator function of the Sphecidae."

**Olaf Zimmermann** (Journalistenbüro Zimmermann & Franken GBR, Lürripen Str. 228, D-41065 Mönchengladbach, Germany), an author, is writing a book entitled "Die sozialen Faltenwespen (Vespoidea)."



### HELP NEEDED

Work Progress on Tachysphex pechumani (Sphecidae: Larrinae) by Mark O'Brien & Frank Kurczewski Museum of Zoology, University of Michigan, Ann Arbor, MI 48109-1079 and Environmental & Forest Biology, SUNY College of Environmental Science & Forestry, Syracuse, NY 13210-2778

Over the last decade, we have been studying the behavior, ecology, and distribution of Tachysphex pechumani Krombein. During the past 3 years, we have greatly increased the number of observations, have added many new localities to its distribution, and have travelled several thousand miles searching for new sites. We are concentrating our efforts in Michigan for, until this summer, the only known populations were situated in New Jersey and Michigan. An extension of the Michigan population was discovered this summer in northwestern Ohio (Lucas Co.), which was a nice range expansion. The New Jersey population had been studied previously (Kurczewski & Elliott 1978), and we are comparing it with the Michigan population.

One of our remaining problems is to fully define the current and past distribution of this wasp species. Although we have scoured collections, written letters to innumerable curators, and searched in other areas where the species might possibly occur, we have yet to find any other states besides MI, NJ, & OH where *T. pechumani* occurs. We plan to go to Wisconsin in the summer of 1994 to search in suitable habitats i.e., sandy soil with oak-pine barrens, savannas and woodlands or compacted areas of open sand (such as two-tracks).

We are asking collectors to examine their collections for *T. pechumani*. It is the only medium-sized *Tachysphex* with orange antennal segments and anterior half of the abdomen red in the female. It is otherwise black, including the much smaller males which are all black. If you know of any suitable habitat in the Great Lakes region that we can examine (IN, OH, WI, MI, NY, SW Ontario?), please let us know. The emergence period for these wasps is early June to mid-late July.

The best way to contact us is to send an e-mail message: (userlps5@um.cc .umich.edu), or call Frank at 315-470-6753, or Mark at 313-747-2199. We will appreciate any leads that our colleagues can offer!

#### References

Kurczewski, F. E., & N. B. Elliott. 1978. Nesting behavior and ecology of *Tachysphex pechumani* Krombein (Hymenoptera: Sphecidae). J. Kansas Ent. Soc., 51:765-780.2



# **NEW ADDRESSES**

- John C. Abbott: 1030 Dallas Drive, Denton, Texas 76205.
- Celso Oliveira Azevedo: Universidad Federal do Espirito Santo, Dept. Biologia, Av. Marechal Campos 1468, Maruipe, 29040-090 Vitória, ES, Brasil.
- David L. Evans: 2616 Lincoln Drive, Montoursville, PA 17754.
- Jeremy Field: Dept. of Ecology & Evolutionary Biology, Rice University, PO Box 1892, Houston, Texas 77251.

- Jorge E. Frana: E. Oliber 1026, (2300) Rafaela, Santa Fe, Argentina.
- Eric Hammarström: C/o Johansson Majstångsgatan 11A<sup>III</sup>, S-414 72 Göteborg, Sweden.
- Dr. Alois Kofler: Meranerstaße 3, A-9900 Lienz /Osttirol, Austria.
- Gabriel Augusto R. de Melo: Snow Entomological Museum, Snow Hall, The University of Kansas, Lawrence, KS 66045-2106.
- Sean O'Donnell: Dept. of Entomology, Univ. of California, Davis CA 95616-8584.
- Michael Ohl: II Zoologisches Institut und Museum, Berliner Str. 28, 37073 Göttingen, Germany.
- Dr. Margery G. Spofford: 112 Seneca Road East, Trumansburg, New York 11886.
- Dr. R.J. Paxton: Department of Genetics, Uppsala University, Box 7003, S-750 07 Uppsala, Sweden (until 30 April 1994).
- Dr. Peter F. Yeo: The Store House, Gretton Road, Harringworth, Northants NN17 3AD, UK.

# **E-MAIL ADDRESSES**

Celso Oliveira Azevedo, Brasil: Celso@Brufes.bitnet Wojciech J. Pulawski, San Francisco, CA : WPulawski@calacademy.org



## **PEOPLE IN THE NEWS**

Julio Antonio Genaro (Apartado 6099, Habana, Cuba) is working at the Museo Nacional de Historia Natural (Capitolio Nacional, Ciudad de La Habana 10200, Cuba) as department chief of collections.

Roy Snelling (Dept. of Entomology, Natural History Museum of Los Angeles County, 900 Exposition Blvd., Los Angeles, CA 90007) comments: "Well, guys, it finally happened, though not on the schedule I had hoped for. But, thanks to the concerted efforts the administration here at good ole LACM, I was forced into an untenable situation and therefore chose to retire. Effective 1 September 1993. As everybody knows by now, the economic situation in the Golden State can only be described as a disaster, precipitated by too many State and Federal administrations convinced that what's good for them and their buddies is more relevant than what's good for the people at large. (Sounds sorta like what happened here on a smaller, more personal scale.) Whatever. LACM took a terrible series of 'hits' in personnel reductions, first in January, then again in August. Very nearly all County positions were eliminated; most of these were 'rolled over' to the LACM Foundation. With one minor exception, all these cuts came from areas other than Museum administration. Why is this no surprise? The Foundation has suffered from serious fiscal mismanagement and I think that a lot more people will be lost from here before all is said and done. The option to return under contract to the Foundation was offered me, but on terms that I found morally repugnant.

"Over the last few years the research and collections component of the Museum has been seriously compromised, with some sections, such as Botany, entirely eliminated. On the more positive side, our current Director will retire in 6-8 mos. and we may be fortunate enough to acquire a new Director who will be better able to manage this place. I do continue to be optimistic that once the present national economic crisis is over, the LACM can be rebuilt to its former status as an institution of some importance.

"I am now a Research Associate, so will continue to come to the Museum with fair regularity. I will, finally, be able to conduct my research with a minimum of disturbance, thereby increasing the liklihood that some of my numerous projects will actually see completion. I admit that it is sort of neat to be able to say to myself, when the alarm goes off at 0430: 'I don't have to go to work if I don't wish to.' Then I get up and go to work."



Cephalopamops denticulatus Spinola, male. (Chrysididae, North Africa, Middle East)

Frank Parker retired from his USDA job in Costa Rica at the end of September. He will spend his retirement in Nevada or Utah. We will publish his new address when available.

# **MISSING PERSONS**

Dr. E.O. Grechka of Kherson, Ukraine Dr. Henry Hermann of Athens, Georgia Dr. Daniel T. Jennings of Orono, Maine Mark Parrish of New Brunswick, New Jersev.

Dr. Tadashi Suzuki of Tokyo, Japan.



# HYMENOPTERA RESEARCH

# Coup Attempt at J. Hym. Res.

An insidious plot was uncovered to turn the latest issue of the *Journal of Hymenoptera Research* into an American publication. On the spine was printed "AMERICAN JOURNAL HYMENOPTE-RA RESEARCH VOLUME 2, NUMBER 1, 1993." Quick thinking and some deft work with black markers by David R. Smith, Arnold and Nancy Menke, with the approval of editor Paul Marsh, saved the day. We would like to assure President-Elect Donald Quicke of Sheffield, England, that his office is secure.

Volume 2, number 1, was mailed November 17, 1993 and that's the correct publication date for this issue. The September 30 date printed on the inside front cover is erroneus.



Nesogyne taino Krombein, female. (Chrysididae: Amiseginae, Dominican Republic

# **BIOTA**

**BIOTA** (Biosystematic Information on Terrestrial Arthropods) was developed as a project to create a nomenclatorial database of the insects of the world. It has since assumed a broader meaning and now serves as an umbrella to cover all of the computer-related information resources existing or under development within the Systematic Entomology Laboratory (SEL) of the United States Department of Agriculture. Information tools underway in the SEL are catalogs (Biosystematic Database of World Diptera, systematic catalog of worldwide cutworm moths), specimen label databases, the Fruit Fly Expert System (automated identification tool for extension specialists and guarantine inspectors), and other automated identification systems. We envision the day when a global network of information databases and other tools developed by a union of individuals and institutions (including SEL) will exist and when anyone, anywhere will be able to access all available information on a terrestrial arthropod species from his/her computer.

The first project of *BIOTA* is a nomenclatorial database. A nomenclatorial database has many uses, among them are the following:

1. A nomenclatorial database is a compendium of the known species of a group, literally a phone directory of the biological sciences. It is the first step in any biosystematic or biodiversity program.

2. A nomenclatorial database can be used to investigate fundamental questions of biodiversity. For example, how are species distributed among the biogeographical provinces? Are predator or herbivore groups more diverse?

 A nomenclatorial database provides stability for names and serves as an authority list for users of biosystematic names.

4. A nomenclatorial database is the first component of any comprehensive biosystematic information system on insects.

5. A nomenclatorial database has vital information for anyone doing biosystematic work on a group of organisms, be it revisionary, curatorial, or biodiversity.

6. The nomenclatorial database in *BIO*-*TA* provides access to the literature by giving the source of the name (revision, catalog, check list, unpublished data) and an authority who may be contacted to provide further information. A current nomenclatorial database of all the scientific names (species-, genus-, and family-group names, including synonyms, homonyms, misspellings, and misidentifications) applied to insects and other terrestrial arthropods has the highest priority because it is the backbone of all information services. The database is an SEL-wide project and is under the supervision of Ronald W. Hodges (chair), Robert W. Poole, and F. Christian Thompson.

The *BIOTA* nomenclatorial database is divided into four data tables: species, genera, families, and ranges (biogeographic regions). A data entry program has been developed for the use of specialists and other contributors. A manual, which provides full documentation on the check list, accompanies the program.

Each individual contributing to the database and his/her institutional affiliation will be fully acknowledged. First, there will be an introductory acknowledgment screen in the finished product. Second, each record of each database contains a memo field. If data are based on the unpublished resources of a contributor, this fact will be noted.

The first two years of the nomenclatorial database project will be spent assembling all of the names for the North American fauna. Data entry will not be restricted to North American species because this is a global project; however, the emphasis will be on the North American fauna. A date for completion of the check list on a world level cannot be estimated accurately because it depends on the collaboration and cooperation of our colleagues throughout the world. Similarly, no accurate estimates of the final size of the project are available. Probably, well over two million names are involved.

The database will be available on the Internet and on CD-ROM.

Contact Ronald W. Hodges for further information.

#### **Ronald W. Hodges**

Systematic Entomology Laboratory, ARS, USDA c/o National Museum of Natural History, NHB-168 Washington, DC 20560 USA (202) 382-1778; FAX (202) 786-9422



# FORUM

# Subgenera and Species Groups

*by* Jean Leclercq

Faculté d'Agronomie de Gembloux, rue de Bois-de-Breux, 190, B 4020, Liege - Jupille, Belgium.

I do not agree at all with the view [See Sphecos 10:11-13, 11:11-13, etc., for discussion of this topic - editor] that subgenera are best re-placed by species groups. I believe that subgenera are needed to indicate an evolutionary (or cladistic) level higher than species groups and I really fail to see that nomenclature is simplified if instead of using subgenera in which there is one name and one author, imposed by the law of priority, we have longer names such as "Ectemnius lapidarius group". (In that case why lapidarius and not sexcinctus, and why not the name of the species which finally will appear to be either the commonest, the best known, or the obviously most primitive?)1

In any case, I think that all formal names in a classification are hypotheses to be tested and tested again, that the species level is the only one likely to represent a final, duly tested and retested, natural category. Subgenera, genera and other higher categories express something, certainly, but not always the same evolutionary reality; indeed they are subjective, pragmatic taxa: they are drawers which must have a name and a coherent content, but which for practical reasons, should not be too large.

Although I believe that subgenera mean something and are to be used in certain projects, e.g. when there are reasons to make the nearest species more easily compared, I am against the use of subgenera in many pratical circumstances:

(a) on determination labels!

(b) in collections – here I favor the alphabetical order of the genera within a subfamily and of species within a genus. The purpose of collections is not to express or suggest taxonomic relationships. When searching in a collection, the curator or the specialist needs to easily determine if a particular species is represented, not to check its place among its nearest relatives.

(c) in files, computerized or not, in atlases of geographic distributions, and other practical documents.

So there is no contradiction in the fact that in the Faune de France - Sphecidae, with J. Bitsch, the species are commented, grouped under their subgenera (this is a comprehensive treatise in which e.g. Ectemnius borealis, dives and guttatus must come together because they are related (in the same subgenus Ectemnius), with information which must be readily compared) and the fact that species are simply in alphabetical order in my list "Hymenopteres Sphecides Crabroniens d' Europe et du Bassin Mediterraneen" (N.F.Gx nº 26, 1993) and in our set of maps J. Leclercq & Y. Barbier "Atlas de repartition des Crabroniens de France et des regions limitrophes" (N.F.Gx, nº 27, 1993).

#### On Identification Keys by Jean Leclercq

[The following was prompted in part by material in Sphecos 19:18 and 20: 6-7 = editor.]

I would like to recount my recent experiences with the keys of R.M. Bohart to the North american species of Crabro (1976), Ectemnius (1979) and Rhopalum (1974). They proved to be concise keys near to perfection. However I think that a user would make 5 to 10% of his determinations wrong and 5 to 10% or more determinations with a "?" if he were a hymenopterist without a reference collection, because there are not enough characters in these keys to verify his results for every species, not enough even if he continued reading what is given further on in the paper. Furthermore, if while using these keys he wanted to know whether one particular Mexican or South American specimen belongs to one of those North American species in the key, well again, there is no answer without a good reference collection.

My experience with the keys of Kohl for Crabronini and other Sphecidae and of Blüthgen for Vespoidea leads me to conclude that these works guarantee more certainty of a correct identification, or more reasons to believe you are seeing another species, because more characters are given, implying that the authors did not so easily think they had selected the most "essential" characters of value for the practical separation of taxa.

In my own work, I often produce keys with more characters than necessary, or wanted by the user. But that is because I deal with exotic faunas, and usually with divergent specimens and small series: careful examination of these specimens does not reveal the characters which would make it easiest to recognise the relevant taxa. Very often I am not sure I am dealing with a true species (named or to be named) but I feel it is important to present my hypothesis of the status of these species (or subspecies) in order to encourage further research by others. So in practice, my keys are a practical system to present the results of my comparisons and to help future taxonomists check their material against my experience, eventually producing improved keys due to better characters. Thus, for exotic material (of course not the European fauna), my keys are not built to permit the easiest determinations by anybody who thinks that should already be possible. Colleagues in a hurry to find a name must be discouraged but those who will intend to do better taxonomic work with new material should, I hope, appreciate that I leave to them a record of what I found and of the problems I faced.

## ONE MAN'S WISDOM ON TAXONOMIC CONTROVERSY

"My personal preference is to keep disagreements polite and respectful, just in case the person with whom I disagree turns out to be right after all."

ex litteris Byron Alexander



Haplogonotopus sp. male (Dryinidae), Australia

The generic name need not be part of the species group name, *lapidarius* group is sufficient. As far as which species name to use, the type-species of the subgeneric name seems a logical choice – Editor.

# PET PEEVE

#### Serving the User

I wish to support strongly Arnold Menke's "Pet Peeve" in Sphecos 25:7. Workers who fail to add identification labels to every specimen they have named are simply wasting their initial investment of time - they are not communicating. They should remember that "No man is an Island": the human species, being a social animal, makes progress by cooperating - that includes communicating. Entomologists (and other authors) divide into two classes, those who publish for themselves and those who publish for others. The first should not publish at all. The second category are distinguished (in both senses) by producing papers which are straightforward for others to understand and use; when they have to decide whether to include or exclude an item, or on a particular way to set out something, the decision is always taken with the user in mind. Ask yourself, "Would I find my own publication convenient to use?" If this aim is not consciously borne in mind, then something less than good communication will result. One small but significant instance happened a while ago: a reviewer of a short paper I had submitted suggested that I give only the author and year of works cited in textreferences. In my reply to the editor, I strongly resisted this. If you give author, year and page, it makes it so much easier to find the reference. How often have you either spent a long time trying to find the part quoted, or given up on finding it altogether, simply because of lack of a page number which was instantly available to the author himself? This is sheer lack of consideration for the user. Incidentally, many editors will not accept taxonomic works without identification keys - if you cannot understand their reasons, you should not be publishing!

> Colin Vardy Dept. of Entomology The Natural History Museum Cromwell Road London SW7 5BD UK

# SCIENTIFIC NOTES

# Passaloecus pictus in Brasil (Sphecidae)

*by* Sérvio Amarante, Museu de Zoologia da Universidade de São Paulo, Caixa Postal 7172.

01064.970 São Paulo, Brasil and

David Vincent, Bee Research Laboratory (PSI), USDA, ARS, Bidg. 476, BARC-East, 10300 Baltimore Ave., Beltsville, MD 20705-2350.

In the New World the sphecid genus *Passaloecus* is restricted to North America (Vincent, 1979). No species are known in the Neotropical Region, the genus being replaced there by *Polemistus*. Amarante collected specimens of a *Passaloecus* in São Paulo, Brasil, and sent them to Arnold Menke for identification. He tentatively identified them as *pictus* Ribaut, a European species. Menke forwarded the specimens to Vincent who confirmed that they were indeed *pictus*.

In the Old World, Merisuo (1974) recorded *pictus* from southern France, the Crimea Peninsula, and Yugoslavia. Dollfuss (1991) listed the species from southern Europe and Turkey. When Ribaut (1952) described *pictus*, he stated that the species nested in the soil in southern France, near Toulouse. Janvier (1961) confirmed that this wasp nests in the ground, and he provided a detailed description of the nesting behavior of *pictus* in France. Janvier found large aggregations of the wasp nesting in sand banks at an abandoned sand quarry of the Fertes-Alais.

Four females of pictus were collected January and November 1989 and January 1990, and one male was taken December, 1991 in Parque da Independência in the Ipiranga district of the city of São Paulo. The grounds of the Parque include some French gardens designed to copy those of the Louvre, and the Museu Paulista (a historical museum), including its former Horto Botanico. The Horto Botanico is now nothing more than a wooded area and this is where the wasps were collected. Hermann Lüderwaldt, one of Brasil's first entomologists, was formerly employed as a gardener in the Museu Paulista. He planted representatives of the main vegetation zones of Brasil: Cerrado, Caatinga, Atlantic Rain Forest, Amazon Rain Forest, and so on in the Horto (Lüderwaldt, 1918). The Horto Botanico, incidently, is the type locality for many species described by Herman von Ihering, Lüderwaldt, and other entomologists. The French gardens contain some exotic plants, and perhaps *pictus* was introduced with soil of some of them. In an attempt to acquire more material of the wasp and its biology, the Horto and nearby French gardens are being searched for likely nesting areas of sand banks and slopes. Trap nests have been set out as well in case the nest building of *pictus* in the neotropics is not limited to soil.

- Dollfuss, H., 1991. Bestimmungsschlussel der Grabwespen Nord- und Zentraleuropas. Stapfia, Publikation der Botanischen Arbeitsgemeinschaft am O. O. landesmusem Linz, 24:1-247.
- Janvier, H., 1961. Recherches sur les Hymenopteres nidificants aphidivores. III. Le genre *Passsaloecus*. Ann. Sci. Nat., Zool. 12:847-883.
- Lüderwaldt, H., 1918. O Herbário e o Horto Botanico do Museu Paulista. Rev. Mus. Paulista 10:289-311.
- Merisuo, A. K., 1974. Zur Kenntnis der europaischen Arten der Gattung *Passaloecus* Shuckard. Ann. Ent. Fennici 40:10-15.
- Ribaut, H., 1952. Especes francaises du genre *Passaloecus*. Bull. Soc. Ent. France 57:23-28.
- Vincent, D., 1979. A revision of the genus *Passaloecus* (Hymenoptera: Sphecidae) in America North of Mexico. Wasmann J. Biol. 36:127-198.

#### Crabroninae vs Larrinae (Sphecidae) *by* Arnold Menke

Usage of family-group names is governed by priority just as with generic and species level names. When Bohart and Menke (1976) published their book, Sphecid Wasps of the World, subfamily names were not studied to see if those in use had priority. In the newly published Hymenoptera of the World edited by Goulet and Huber (1993), I examined Bert Finnamore's section on Spheciformes, I noted his use of Crabroninae instead of Larrinae for the assemblage containing both Larrinae and Crabroninae of Bohart and Menke (1976). My curiosity was piqued because there was no explanation for using the subfamily name Crabroninae instead of Larrinae, but I assumed that Bert had discovered that Crabroninae was the older of the two family-group names. Fortunately Handlirsch's treatment of Hymenoptera in Handbuch der Entomologie. volume 3, edited by Schröder (1925) lists most family group names (including synonyms) and their authorship. Thus it provides a good starting place for the origin of many higher category names in Hymenoptera. From Handlirsch's work I ascertained that Latreille authored both names: Crabronites dating from 1802, page 340, and Larrates dating from 1810, p. 438. Thus Finnamore was correct in using Crabroninae instead of Larrinae. I should point out here that Finnamore accepted the views of Evans (1964), Lomholdt (1985) and Menke (1988) in uniting the Crabroninae and Larrinae (sensu Bohart and Menke) to form one subfamily. Of course Finnamore recognized all of the subfamilies of Bohart and Menke as families, but that is another matter (see review on page 17).

Michener (1986) reviewed the familygroup names for bees. In view of the considerable study now being given to the classification of Sphecidae by Alexander, Prentice, Ohl, van Zuijlan, and others, and the possible changes in groupings resulting from their work, it is time that the family-group names be investigated so that priorities will be easy to determine. Woj Pulawski and I have to deal with this problem in connection with our work on a new Catalog of Sphecidae and may, therefore, produce a separate paper on family-group names and their authorship.

#### Literature cited

- Bohart, R. M. and A. S. Menke, 1976. Sphecid wasps of the world, a generic revision. University of California Press, Berkeley. x + 695 p.
- Dupuis, Claude, 1986. Dates de publication de l' "Histoire naturelle générale et particulière des crustacés et des insectes" (1802-1805) par Latreille dans le "Buffon de Sonnini". Ann. Soc. Ent. France 22:205-210.
- Evans, Howard E., 1964. The classification and evolution of digger wasps as suggested by larval characters (Hymenoptera: Sphecoidea). Ent. News 65:225-237
- Goulet, H. and J. Huber, 1993. Hymenoptera of the world: an identification

guide to families. vii + 668 p. Centre for Land and Biological Resources Research, Agriculture Canada, Ottawa.

- Finnamore, A. T., 1993. Series Spheciformes, p. 280-306 *in*: Goulet and Huber Hymenoptera of the world: an identification guide to families. vii + 668 p. Centre for Land and Biological Resources Research, Agriculture Canada, Ottawa.
- Griffin, Francis J., 1938. On the date of publication of Latreille (In Sonnini's Buffon), An X [sic], Hist. Nat. Gén. Partic. Crust. Ins. 3. J. Soc. Bibl. Nat. Hist. 1:157.
- Handlirsch, A., 1925. Geschichte, Literatur, Technik, Paläontologie, Phylogenie, Systematik, Vol. 3, viii + 1202 p. *in*: Schröder, C., Handbuch der Entomologie. Gustav Fischer, Jena.
- Latreille, P. A., 1802. Histoire naturelle, générale et particulière des crustacés et des insectes. Vol. 3. F. Dufart, Paris. xii+13-467 p. Published between September and November 6, 1802 (see Griffin, 1938 and Dupuis 1986).
- Latreille, P. A., 1810. Considerations générales sur l'ordre naturel des animaux composant les classes des crustacés, des arachnides, et des insectes. F. Schoell, Paris. 444 p.
- Lomholdt, O. C., 1985. A reclassification of the larrine tribes with a revision of the Miscophini of southern Africa and Madagascar (Hymenoptera: Sphecidae). Ent. Scand., Suppl. 24, 183 p.
- Menke, A. S., 1988. *Pison* in the New World: a revision (Hymenoptera: Sphecidae: Trypoxylini). Contrib. Amer. Ent. Inst. 24(3):1-171.
- Michener, C. D., 1986. Family-Group names among bees. J. Kansas Ent. Soc. 59:219-234.

#### Larra altamazonica in Panama by Arnold Menke

This species is known from northern South America including western Colombia (Menke, 1992), but *altamazonica* has not been recorded from Central America. In February, 1993, Roberto Cambra collected one female in Panama: Darien: Cruce de Mono, Estacion Inrenare, Parque Nac. Darien (Museo de Invertebrados G. B. Fairchild, Panamá, Panamá). The specimen represents the form with a red abdomen. Literature cited

Menke, A. S., 1992. Mole cricket hunters of the genus *Larra* in the New World, J. Hymen, Res. 1:175-234.

#### Additional records for New World *Pison* by Arnold Menke

#### Pison eu Menke

PANAMÁ: Darien: Cruce de Mono, Estacion Irenare, Parque Nac. Darien (Museo de Invertebrados G. B. Fairchild, Panamá, Panamá), one male.

This specimen has weak tyli on flagellomeres V-VII. The OOD is .9X the HID and the UID is .9X the LID. This is the first record of *eu* from Panama.

#### Pison gnythos Menke

PANAMÁ: same locality data as above, one female.

This is the first record of *gynthos* from Panamá although it is known from Costa Rica and Colombia.

# The Possible Prey of Laphyragogus aljer deBeaumont

*by* **Michael Prentice** Dept. of Entomology, University of California, Berkeley, CA 94720

In Sphecos 10, Kazenas presented brief observations on Laphyragogus turanicus Gussakovskij describing a wasp provisioning with pyralids of the genus Crambus. Previously, nothing had been known of the habits of this peculiar sphecid genus which Bohart and Menke (1976) place in its own subfamily. To Kazenas' observations, I may add that I found many moth scales on a female specimen of Laphyragogus aijer deBeaumont sent to me by Kenneth Guichard. Possibly this species, like turanicus, preys on small Lepidoptera, although the scales may, alternatively, be the result of collecting. Perhaps Guichard knows whether the collector (I. L. Hamer, Oman, April 17, 1992) was likely to place moths and wasps into the same jars (obviously a poor practice). Since aijer is the most distinct of the six species of Laphyragogus, it seems likely that all prey on moths and that the unusual morphology of the genus is in part a specialization for moth collecting.

# Early Cretaceous Aculeata from Lebanese amber by Michael Prentice

With the help of Dr. George Poinar of U. C. Berkeley, I have had the opportunity to examine two interesting collections of Lebanese amber. This amber has the distinction of being the oldest known to contain insect inclusions and is minimally dated to 135-120 Ma (Earliest Cretaceous), an important period in aculeate evolution. The insects preserved are for the most part small (usually 5mm or less), although a few particularly interesting ones exceed this and there are many fragments of larger insects. My impression is that the amber is comparatively rich in Aculeata (seemingly more so than Dominican amber) although I have only been able to look through a fraction of the material to this point.

There are at least 15 aculeates, some just represented by wings. Several of the wasps may be tentatively assigned to extant families including Scolebythidae, Dryinidae(?), Bethylidae and Sphecidae, while others are probably new. The most impressive is a female ampulicine, approximately 4mm in length, which, unfortunately, lacks a portion of the metasoma and most of the wings. I have been able to only briefly examine it, but can report that it is unquestionably an ampulicine, resembling Ampulex to a high degree with an elongate pronotum. However, it lacks the wedgeshaped clypeus and greatly elongate mandibles of female Ampulex. This is the oldest undoubted sphecid and indicates that the family (in the broad sense of Bohart and Menke (1976)) may extend well back into the Jurassic! That it appears so close to an extant genus is incredible, and indicates Ampulicinae have changed little since the beginning of the Cretaceous. In one of the collections there is also a much smaller ampulicine-like wasp, around 1.5mm in length, that i have not had a chance to study in detail.

Among the other finds is a well preserved scolebythid which I am currently describing. It has the characteristic propectal morphology of the family, but is more primitive with respect to the pronotum which is dorsally long, unlike the derived, short pronota of living scolebythids. This feature, by the way, supports Brothers' (1975) and Carpenter's (1986) view that the shortened pronota of Scolebythidae and Plumariidae evolved independently. Another less well preserved wasp has a scolebythid appearance, and a number of disassociated wings have scolebythid-like venation suggesting that the family was considerably more common in the Early Cretaceous than today.

Among the forms that may represent new families is one comparatively large wasp (around 5mm in length). Due to the present state of polish of the piece, it is impossible to provide many details except to say that this wasp appears very primitive, resembling a squat scoliid in outline. The remaining aculeates are in need of extensive polishing, but are all chrysidoid-like in appearance.

When properly studied, these specimens, along with others that will surely be uncovered, will provide an unequaled view of early aculeate evolution.

#### A Social Wasp Which Stores Nectar by Martin Cooper "Hillcrest", Ware Lane, Lyme Regis, Dorset D7T 3EL England

My friend Dr. G. Onore of the Universidad Catolica in Quito tells me that *Polybia a. aequatorialis* Zavattari, a social wasp of the montane zone in Ecuador stores nectar in its nest. At Oyacachi, a village in the province of Imbabura at 3,200m, the country folk relish the combs containing nectar as a delicacy. It is well known that two species of the Neotropical genus *Brachygastra*, *lecheguana* (Latreille) and *mellifica* (Say) also store nectar in their nests.

I examined a nest of *aequatorialis* which Onore had collected. It was a typical *Polybia* nest built around the twig of a shrub: pyriform, 12cm long by 9.5 cm wide with four combs. The envelope was pale brown, 1-2cm wide with pockets forming air spaces between 3-4 layers of carton. The spaces in the envelope probably help to insulate the nest during the cold Andean nights. I found neither nectar nor brood in this nest which had been collected four months previously.

A. aequatorialis in Colombia and Ecuador forms a mimicry group with two other social wasps: Agelaia baezae (Richards) and Mischocyttarus rufomaculatus Richards. This mimicry group is not recorded by Richards (1978 p.30). The ground colouring of the wasps is black except for the anterior half of the gaster which is yellowish or pale reddishbrown. I found all three species in the Banos Vally of Ecuador together with other insects belonging to the same mimicry group. These were: a Cerceris sp. (Sphecidae), a halictine bee, an ichneumonid wasp and a Tipula (Diptera) which flies in a manner reminiscent of a hunting social wasp.

#### Reference

Richards, O.W. 1978. The social wasps of the Americas excluding the Vespinae. British Museum (Natural History), London. 580 p.

#### Defensive Behaviour of *Chartergus chartarius* (Oliver) *by* Martin Cooper

It is well known that wasps of the genus Synoeca make a noise like someone sawing wood when their colonies are alarmed; hence the name "guitarrera" (guitar player) in Columbia. The wasps make the sound by rasping the envelope with their mandibles in unison. Another species of social wasp which makes a sound when alarmed, though less intimidating, is *Chartergus chartarius* (Oliver).

When I once disturbed a nest of this species the workers aggregated on the outside of the envelope with their abdomens raised. As I continued to molest them (I wanted the nest as a trophy) the wasps became more excited and began to beat their abdomens against the envelope to make an audible drumming sound. Such a noise could serve either as a warning to a potential predator or to arouse the workers still inside the nest.

The envelopes of *Chartergus* nests are white and the wasps black so they are very conspicuous when assembled on the envelope. Some nests have black patches, perhaps to simulate an aggregation of wasps. Other species which are black and have nests with white or whitish-grey envelopes are: *Pseudopolybia compressa* (Saussure), *Epipona* sp. and some *Parachartergus*.

#### Unusual Behaviour of Female Agelaia testacea (F.) by Martin Cooper

When in a Bolivian forest a few years ago I came across two Agalaia testacea (F.) behaving in a manner such as I have never seen before in a social wasp. The wasps were repeatedly flying without touching at about two feet from the ground. I have seen a similar confrontation between two male *Trypoxylon* (*Trypargilum*) of a species in the *nitidum* group (Sphecidae) and two males of an unidentified bombiliid fly. However when I caught the two *A. testacea* they turned out to be females.

#### Unusual Mating Behavior by Robert S. Jacobson Dept. of Pathology and Laboratory Medicine, School of Medicine, East Carolina University, Greenville, North Carolina 27858-4354

While collecting Bombus and Polistes from flowers on the afternoon of 24 October 1993 in Beaufort Co., North Carolina I saw what appeared to be an attempt by a male of Bombus fraternus (Smith) to mate with a young queen of Bombus impatiens Cresson. These were together inside my net along with a couple males of impatiens, another male of fraternus, and a male of Bombus pennsylvanicus (Degeer). The rather senescent fraternus male persisted in mounting the female and extruding his claspers. As far as I could see, the female was not receptive and I doubt the male had much success. They remained together even after being placed into a rather weak cyanide jar. Unfortunately, no fraternus queens were seen that afternoon, in order to give this male a choice. I find it amazing that a male from a different subgenus would attempt this mating. Perhaps their being as close together as they were in a net artificially bypassed isolating mechanisms that would otherwise prevent it, but what if these two had landed by chance on the same flower? Comments or questions are welcome. I should add that neither of the impatiens males were observed attempting to mate with this queen, which appears slightly callow.

### Similar Mating Behavior in Ants

I once observed similar behavior in flying ants. One night on the campus of North Carolina State University in Raleigh, many species of ants were having their mating flights. In an aspirator I collected a diverse group including a female Pheidole vinelandica Forel (Myrmecinae) and some male Paratrechina vividula Nylander (Formicinae). Upon finding themselves in close guarters in a collecting bottle, all the male Paratrechina attempted to mate with the female Pheidole dispite being from different subfamilies. I assumed that such unusual behavior was an artifact of being in unnatural conditions. Somehow, being confined short-circuited the normal behavioral sequences that would allow proper descrimination of the correct female species.

Terry Nuhn

# **TECHNIQUES**

#### Yellow Pan, an Easy and Very Effective Way for Catching Spider Wasps.

by Raymond Wahis Faculté des Sciences agronomiques Zoologie générale et appliquée B-5030 GEMBLOUX (Belgium)

Ma fille cadette séjournant pour une année au pair dans une famille américaine, à Alexandria (Virginia), je profitai de cette occasion pour une première visite au USA où je séjournai une douzaine de jours dans sa famille d'accueil (Family Teague, Emerald Drive), dans un environnement résidentiel boisé. à Wavnewood, à proximité immédiate du parc de Fort Hunt. Derrière l'habitation, la pelouse donne sur une parcelle boisée avec de hauts arbres tels: Liriodendron tulipifera, Liquidambar styraciflua, Sassafras albidum, Robinia pseudoacacia, Hamamelis virginiana, Acer nigrum, Fagus grandifolia, Quercus alba et coccinea, etc. La strate arbustive se compose principalement de Comus florida et de rares llex opaca. Le sous-bois, à sol en majeure partie couvert d'une litière de feuilles mortes. est relativement sombre avec toutefois quelques spots ensoleillés peu étendus, se déplaçant au cours de la journée suivant la position du soleil.

Ayant aperçu et capturé, sur la pe-

louse, un Tachypompilus ferrugineus 9 trainant sa volumineuse proie (Dolomedes tenebrosus Hentz, Pisauridae, det. Coddington) en direction d'un mur de l'habitation (preuve de la présence de Pompilides), je disposai à la lisière de la forêt et dans le sous-bois une vingtaine de bacs en aluminium léger (20 cm x 13 x 5), peints en jaune clair et remplis d'eau additionnée de quelques gouttes de détergent (matériel utilisé en alimentation pour la présentation des hors-d'oeuvres et plats préparés). Ces bacs restèrent en place pendant 6 jours seulement, du 9 au 14.VIII.93 (laps de temps, que d'expérience, i'estime largement suffisant pour une première analyse). Chaque soir, je prélevais les hyménoptères piégés pendant la journée, nettoyais les bacs et rajoutais, si nécessaire, un peu d'eau pour compenser la perte par évaporation.

Ce système est particulièrement efficace pour les Pompilides généralement considérés, avec raison, comme de capture difficile. Il permet notamment la prise d'espèces de petite taille lesquelles, se déplaçant dans la végétation ou au sol dans la litière de feuilles mortes, passeraient autrement inaperçues. Dans ce milieu apparemment peu propice, j'ai pu ainsi récolter sans peine 51 spécimens répartis en 15 espèces.

Je ne saurais donc trop recommander l'utilisation de ces bacs, peu coûteux et faciles à placer. Moins encombrants et plus discrets que d'autres pièges, comme le MALAISE, leur emploi, voire en complément de la chasse traditionnelle au filet, permet, sans y consacrer trop de temps et d'efforts, de dresser un inventaire relativement complet des espèces les plus représentatives d'un milieu donné.

Pepsinae:

Priocnessus nebulosus (Dahlbom) 3 ♀ Priocnemis (s.str.) scitula scitula

(Cresson) 1 9

- Caliadurgus fasciatellus alienatus (Smith) 2 9
- Dipogon (s.str.) brevis recalvus Townes 2 9

Dipogon (Deuteragenia) papago anomalus (Dreisbach) 1 9

Dipogon (Deuteragenia) sayi sayi Banks 2 S

Auplopus mellipes mellipes (Say) 5 Auplopus nigrellus (Banks) 1 Ageniella (s.str.) norota Banks 10 Pompilinae:

Tachypompilus ferrugineus ferrugineus (Say) 1 ♀ avec sa proie, Dolomedes tenebrosus Hentz [det. Coddington] Episyron biguttatus biguttatus (Fabricius) 2 ♀ 1 ♂ Evagetes parvus (Cresson) 2 ♀ Aporus (s.str.) niger (Cresson) 2 ♀ Allaporus pulchellus (Banks) 12 ♀ 2 ♂

#### Ceropalinae:

Ceropales hatoda Brimley 2 9 1 ð

### Chloroform Gun Again *by* Colin Vardy

Dept. of Entomology, The Natural History Museum, Cromwell Road, London SW7 5BD UK

Servio Amarante's account (Sphecos 25:16-20) of his collecting trips is interesting, but he's apparently never heard of the "Chloroform Gun" technique! I think Arnold Menke just had bad luck with trying it (on Ampulex, see Sphecos 18:17-19). I also failed once (or almost so) with Sphecius in Argentina. These two genera must have some kind of armour-plated resistance; it doesn't work on tarantulas, either. But I can assure you it worked very well with all the other wasps (and bees) I tried it on even giant Pepsis in flight "crashed" in only 2-3 seconds; I got them as they rose from a muddy pool. Great sport!



Haplogonotopus americanus female (Dryinidae).

# **COLLECTING REPORTS**

# Egypt Revisited

Wojciech J. Pulawski Dept. of Entomology, California Academy of Sciences, Golden Gate Park, San Francisco, CA 94118

My first acquaintance with Egypt was back in 1957/58, when I spent seven and half months in that country owing to a cultural exchange agreement with Poland. It was a tremendous experience which influenced the rest of my life. All these years I remembered the proverb "elli ishrab maya el Nil, yerga tani". For those who are not fluent in Arabic, this means "whoever drank water of the Nile, will come back again." Indeed, I came back in April 1993, as a leader of a tour group sponsored by the California Academy of Sciences, my mother institution now. When the tour (7-20 April) was over, I stayed an additional six weeks wasping, leaving 2 June.

Egypt has changed dramatically during these years. The population has doubled, Cairo has become a 15 million monster (up from some 4 million people in the old days); roads have been upgraded and new ones built; the old oxen or camel driven sakiahs (irrigation wheels) have almost disappeared and been replaced by modern pumps; new houses and particularly new, pretty looking mosques can be seen everywhere. Yet, many things remain the same: the canals, the fellahin, the crops, the ruins, the sun, and the unique sense of etemity.

The demographic explosion and overall progress have greatly affected nature. The classic collecting sites near Cairo, the type localities for hundreds of species, are nearly all gone. In the old days, I collected insects on the desert outskirts near the Ghiza pyramids, at Abu Roash, and Maadi. All these places are now under construction, or are covered with industrial and house rubbish. Sad indeed. The road from Cairo to Alexandria that I remembered as running through the desert is now boordered by plantations and even new cities (such as Medina el Sadat). The most dramatic changes can perhaps be seen along the Mediterranean coast between Alexandria and Marsa Matruh. Some 200 kilometers are almost entirely covered by new resorts, on a massive scale. An

extraordinary engineering, construction, and overall planning achievement, but what habitat destruction! The uncontrolled population growth (about 2% a vear) is doubtless the most serious problem that the country is facing. For example, for the first time in its history, Equpt became a food importer, and most people have to wait years in order to get a new apartment in spite of all the government efforts. New jobs can barely keep up with the number of new babies. The standard of living just cannot be raised as long as most women continue to became pregnant every 10 months.

One question that I was frequently asked before the trip concerned the muslim fundamentalists, following their attacks on tourists back in 1992. The danger was grossly exaggerated, indeed blown out of proportion by the press and television. Although our group was escorted by police in Cairo, I never encountered an unfriendly gesture, either while being part of the tour, or traveling on my own. People depend so much on tourism that any decline in visitor numbers badly affects merchants, guides, producers of souvenirs, feluca and camel owners, hotel and restaurants owners, taxi drivers, literally millions of people. Besides, Egyptians are just naturally friendly. True, one is constantly exposed to commercial harassment in all touristy places: "Come and see my shop," "Do you want a camel?" "A feluca. Messiew, a feluca," "Do you need a guide?", "Where are you going?", "What country are you from?", "What's your name?", "Taxi, taxi!", "Give me a bakshish!", to give just a few examples. Since I learned some rudimentary Arabic, I was able to demonstrate that I was not a novice and did not need all these services. After landing on the west bank in Aswan, I was immediately approached by a group of camel drivers vehemently offering their services. I just explained that I was not a tourist, showed my net and killing iars, and I how I catch insects. All the boys were greatly interested. When 1 asked if I could go into the gardens, the answer was: "You can go everywhere." I was never molested again there that day or any of the subsequent days. I used this strategy everywhere, especially when collecting near someone's field or garden. After all, they were entitled to know what I was doing there, and ignoring them would only create justified animosity. Many places in Egypt are incredibly cheap these days. Our group stayed in a Hilton hotel in Luxor, but I moved out as soon as I was on my own (because of its limited resources, California Academy of Sciences is not renowned for paying high salaries or being overly generous in financing fieldwork). I found a hotel with air conditioning and a separate bathroom with shower that charged 15 Egyptian pounds per day, or US \$5. It was simply a little deeper inside the local section of the town, just a 15 minutes walk away from the temple and the Nile. One can stay in Cairo for the same price, although these places are rather drab. Hotels in Sinai or along the Mediterranean coast, however, are markedly more expensive.

Collecting in Egypt is notoriously difficult. On one hand, most of its territory is desert, a large part of which is absolutely sterile, with no trace of life. The Nile valley, on the other hand, has been turned into fields and gardens since pharaonic times. In addition, the valley was annually inundated until the first Aswan dam was constructed early this century. The best chance for an insect collector is the narrow contact belt between the desert and the cultivated zone. Another good possibility for insect life is offered by numerous wadis, or arroyos as they are known in North America. Several of them became accessible recently owing to new roads.

Well, when the group left, I stayed in Luxor for another week and went back immediately to my old collecting site on the west bank, no more than a hundred meters away from the spectacular Medinet Habu temple. Yes, it was there, an unspoiled garden, even better than before. Wasps and bees were swarming, both on the ground and on parsley flowers. Many found their way to San Francisco.

A week in Aswan was another success. Although most areas are either fields or human settlements or just a sterile, stony desert, the west bank offers possibilities for wasping. There are some small gardens, several of which were unused and thus full of insects, and further south you find steep cliffs and sand dunes. Good habitats could be found just along the water. And the water is still clean: I swam in it, a refreshing experience during the oppressive heat. I could see fish, plenty of algae, many herons and all kinds of other birds. Amazingly, there were no people there, and I was completely alone, just facing Kitchener Island, one of the famous tourist attractions.

I would like to mention three other places in Upper Eqypt. One is Abydos, the birthplace of Osiris and a famous archaeological site. I went to El Balyana by the 6 AM train from Luxor, and took a communal taxi from there for a ridiculous price of US \$0.30 or so (the distance is about 10km). We ended near the temple, and then I walked through flat sands toward the steep limestone cliff and the canyon that I remembered from the old days. I reached the canvon's mouth in about an hour and indeed, here it was again. Another half an hour up the canyon, perhaps, and I found a blossoming bush full of sphecids and pompilids. I had a badly upset stomach that day (the only time during the trip), and had just enough strength to sit down under a vertical canyon wall, in the shade, to watch the insects fly and to hear a pair of ravens crowing over me. The silence was absolute, the air pure, the sand and rocks unpolluted. I regenerated soon and did my work. And what a pleasure it was to find, higher up, the old sand dune that crossed the canvon and functioned as a natural dam! It was about my own height back in 1958, but now up to perhaps 60 meters. I went over it and found, on the other side, hundreds of small black, dead bushes, but two or three large ones were still alive. What a catch that day!

Another trip from Luxor took me to Qena-Safaga road. This is a recently constructed road that goes from the Nile valley to the Red Sea. I had observed it from the air-conditioned bus, in which our group traveled from the Red Sea to Luxor. We stopped at a small restaurant in the middle of a lush wadi and I knew I would return there again for collecting. When on my own, indeed, I took a taxi to Qena, walked to the Safaga road, and asked the policemen at the road checkpoint, how to go to "kilo hamsa oo tamaneen" (kilometer 85). We expressed all possible greetings first (Salam Aleikum! Aleikum salam u rahman Ullahi wabara katuh! and so on), then one of the men stopped a huge truck and put me on. The driver knew perhaps 10 words of English, my Arabic was only marginally better, but we had a great time conversing all the way to my destination. Collecting was slow but quite valuable. I subsequently discovered that I could go to the place directly by bus from Luxor.

Kom Ombo is only about 50km north of Aswan, with its spectacular Ptolomaic temple facing the Nile. The old gardens that were full of insects between the temple and the shore have been all replaced by pavement. Yet, abandoned rice fields east of the temple turned into nesting sites for aculeates and the trip was not a waste of time.

Finally I flew to Cairo and met Signor Alessandro Mochi of Rome, with whom I previously traveled to the lvory Coast and Senegal. He was born in Cairo and speaks Arabic in addition to English, French, German, and his native Italian, He knows the country very well and was an invaluable guide. I should remind the Sphecos readership that Mr. Mochi owns an excellent collection of Sphecidae (see Sphecos 14:42-44 for details). Most of the original collection, which was built by his father in the 1930s, ended elsewhere as a consequence of the war (in the A. Alfieri collection in Cairo, from where many types and some other specimens were transferred to the USNM, Washington). Nevertheless, Sandro has been able to increase its size significantly through extensive collecting in various parts of Africa during his employment with the World Health Organization, and particularly in Egypt since his retirement several years ago. One difference between us is that he has never been stung by Miscophus mochii, but I was during our expedition. In any case, I greatly enjoyed Sandro's company and I only hope that my feelings were reciprocated. We chatted over various subjects and I would like to record one conversation that we had in Marsa Matruh:

Him: "How many specimens did you collect today?"

Me: "I do not know".

Him: "What would Arnold Menke say?"

Me: "Ignorance is no excuse."

Sandro and I rented a car in Cairo (there are no car rental agencies in Luxor or Aswan, hence I had to rely on public transportation there). We went to Sinai, from Suez to the southern tip of the peninsula and then north again. We explored the area east of Mitla Pass, went to the famous St. Catherine monastery, and almost made it to El Arish on the north coast. We were stopped at a military post (the road was closed to the civilian traffic, a fact not indicated on the map). We had to drive all the way back to Ras Soudr on the west coast, but it was worth it. The following two days we discovered new places with flowering bushes that were loaded with wasps, especially in Wadi Gharandal. Remember this name if you ever go to Sinai.

The last part of our trip was to the Mediterranean coast of Egypt. We passed the World Word II cemeteries at El Alamein and stopped at Marsa Matrouh. What a contrast! The wind from the sea was cold, and it was chilly at night. I saw many plants that I remembered from southern Europe, particularly various composites. Here again we were quite successful, prospecting the area west of the town. The best sites were areas covered by a prostrate *Polygonum* in full blossom, teaming with species that we had not found in other parts of Egypt.

Our last day was spent at Karanis in the Fayum Province, some 65km south of the Ghiza pyramids and next to Greco-Roman ruins. Sandro knew of several *Tamarix* bushes just on the sands edge. They were excellent for wasps and bees indeed. We came back to Cairo that afternoon, returned the car to the renting agency, and I was flying back to London the following morning. Shall I ever see the pyramids again?

Unfortunately, we were not able to meet Giles Roche, who has actively collected around Cairo in the past eight years, discovering excellent new collecting sites (such as Karanis or Wadi Hagul). He had just left Egypt by the time we were there, but his experience was not lost: he had shared his knowledge with Sandro in previous years, and Sandro passed it on to me.

As a whole, the trip was quite a success in spite of the initial disappointment near Cairo. Studies on the Egyptian Sphecidae started more than a century ago (Spinola, 1839), and yet species previously unknown from the country are constantly being added. This trip brought several such additions. I was able to solve some taxonomic problems in *Tachysphex*, although new ones came to light. The most interesting discovery was an unusual prey of a *Gastrosericus*, about which I will give details in my monograph of the genus to be published soon.



#### Collecting Hymenoptera in Southern Maine by Monica Russo 1 North Skilling Road, RR4 Arundel, Biddeford, Maine 04005

On August 13, I was invited to attend a summer field trip organized by the University of Maine at Orono. Entomologists from the state Insect & Disease Lab, the Maine Department of Agriculture, and the Cooperative Extension Office Pest Survey Group participated. The collecting/survey site was the Kennebunk Plains, an extensive area of sandy, open, low-bush blueberry fields along Route 99. There are about one thousand acres of blueberry, scrub oak, shrubby pitch pines and American hazelnut bushes. The land is managed and protected by the Nature Conservancy and is well known locally as a birder's hot-spot.

The expedition was not actually focussed on Hymenoptera at all, but there were a few highlights that **Sphecos** readers might be interested in. At least three *Pelecinus polyturator* females were found. Two of these were easily approachable, user-friendly specimens which I could hand-pick from the vegetation and inspect with an Agfa loupe. Magnification (8X) of the thoracic surface showed intricate sculpturing and textures – a tough challenge to any illustrator looking for a great black-andwhite study subject!

Sandy roads twine across the plains, so we caravanned from one site to another, getting out to scout along the dry dusty lanes. I had the fortune to be standing puzzled by a group of small holes at the edge of the road, when a hefty great golden digger wasp (*Sphex ichneumoneus*) flew up, and I netted her. These diggers are recorded (see reference) from three other sites in Maine, and also at the Maine Audubon sanctuary, but nevertheless it's a real treat to observe one of these colorful, industrious wasps.

About thirty feet farther down the road, another digger was struggling on a bush and then glided down to a hole right in the center of the road. She'd had some trouble flying because she was bringing in a katydid about as long as her own body. She left the katydid carcass at the entrance, and immediately crawled down the hole. By this time everyone had gathered around to watch her efforts. She re-appeared shortly, dragged the katydid down out of sight, and for about thirty seconds we could hear her buzzing exuberantly in her burrow, loudly enough for those standing by to hear her!

I observed several suspected pompilids, crawling and searching the ground with their distinctive where-did-I-dropmy-keys behavior; however, these were not collected or identified that I know of.

One unidentified mutillid was collected early in the day by Don Ouellette from the I&D Lab. This specimen was a rich reddish brown, and the end of the abdomen was black with at least two white lines. At the end of the day when we decided to wrap things up, someone discovered a second mutillid and gave a shout, since Dick Dearborn from I&D was anxious to see a second specimen. The little wasp was in the dirt road, but scurrying rapidly. I helped to corral her by hand, and at one point she even crawled around on my hand, but without any deployment of weaponry. Dick got her into a container, and his day was made!

Field collecting and observing in a unique habitat area like the Kennebunk Plains is a pleasure, especially when one is used to Maine's red maple swamps, and mixed hardwood/softwood secondgrowth forests. I was impressed by the sharing, cooperative nature of all the participants, their enthusiasm and sense of humor, and was grateful to Don for the invitation to attend such a wonderful field trip!

#### Literature Cited

Dearborn, Bradbury & Russell, 1983. Forest Insect Survey of Maine: Order Hymenoptera. Augusta.

#### Back to the Virgin Islands by Roy Snelling Dept. of Entomology, Natural History Museum of Los Angeles County, 900 Exposition Bivd., Los Angeles, CA 90007

I have, of course, been back to the British Virgin Islands. Twice. Mid-April to mid-May, then mid-June to mid-July. The earlier trip began at the end of the dry season and it was dry, indeed. Also damned little in the way of flying Hymenoptera; the creepies (ants) were there as usual. Between the two trips I managed to collect a few more specimens of the new species of Psorthaspis; I've decided to describe the thing myself and I have an illustrator doing a color rendition of the female, a truly gorgeous thing. Also picked up more specimens of the little Pseudomethoca (Mutillidae), including a female. I'm now pretty sure that this is undescribed, but mutillids are someone else's worry, at least those of the Caribbean. Mike lvie sent me some really nifty ones from Hispaniola, most of which seem to be undescribed species. Oddly, there seem to be no records from Puerto Rico. That is surely an artifact of collecting and I'm willing to bet they're there; somebody needs to look in the dry forest areas around Guanica. Females of the Guana Island species are hard to spot. They run about in the leaf litter and at the slightest hint of danger duck into the leaves, never to be seen again. I managed to collect only one; so far, none have turned up in pitfalls.

Incidentally, I must thank Francois Schwartz of Paris. He wrote to advise me that I erred in stating in an earlier Sphecos [24:17] that no mutillids had been previously recorded from the Virgin Islands. There are two, both in *Pseudomethoca: P. olgae* Schuster (St. Croix;  $\Im$  only); *P. unicincta* Ashmead (St. Vincent; both sexes, but the association may or may not be correct – why else would there be an Ashmead Club?). Schuster (1945, Bull. Brook. Ent. Soc. 40:7-8) gives a key to the few thenknown "West Indian" (actually Greater Antillean) species.

To the wasp fauna of Guana Island we can now add *Tiphia* sp. Since the Greater Antillean species of this genus are yet to be worked out, there is little that can be done with it at this time. It doesn't seem to match up with any of the Cuban or Hispaniolan *Tiphia* recorded by Allen & Krombein (1961, Trans. Amer. Ent. Soc. 87:57-66). Only one female.

That reminds me. Here's a PUZZLE-MENTIA for all you brainy types out there. A "how come". How come male and female tiphiids of the same species often have very different wing venation? Since I've never been much of a tiphiid fan, I never really looked at wing venation in the group until I had to make up a key to the families of aculeates found on Mona Island. I made note of a certain venational feature, then when I went back to verify, I happened to pick up a specimen of the opposite sex from the one I'd first examined. Male and female *Myzinum* have very different venation. Weird. I knew there was a good reason I didn't trust tiphilds.

Also to add to the wasp fauna of Guana Island: *Stenodynerus* sp. (Vespidae). Cute, but I don't know what species. And I got a single male of a *Priochilus* (Pompilidae). There is a species of *Priochilus* known from Jamaica, but <u>apparently</u> none known from the Puerto Rico Bank.

Pachodynerus atratus (Fabricius) (Vespidae) was accidentally left off my list (**Sphecos** 24:18); it is common there and was recorded from Guana and various other islands in the area in Menke's paper (1986, Proc. Ent. Soc. Wash. 88: 650-665).

Staying on the subject of Guana (and the V. I. generally), some addenda et corrigenda to my list re the Sphecidae. The wasps that I recorded as *Sphex ichneumoneus* should have been *S. dorsalis* (Fabricius). I goofed.

Tachytes tricinctus (Fabricius). A female collected by Mike Ivie on Saba I. (nr. St. Thomas) was determined by Dick Bohart as *T. distinctus*; looks to be the same as my species and I don't know how they differ or if they do. Fabricius' species was described from the West Indies; *T. distinctus* is a mainland form.

Hoplisoides ater (Gmelin). Bohart determined a specimen from St. Thomas as *H. tricinctus* (Fabricius); I think that its the same as my Guana Island species, probably the correct name, too.

Bicyrtes spinosa (Fabricius). Bohart identified species from St. Thomas for Mike Ivie as *B. discisa* (Taschenberg). J.B. Parker (1929, A generic revision of the fossorial wasps of the tribes Stizini and Bembicini with notes and descriptions of new species; Proc. USNM 75: 1-203) notes that male *discisa* have flagellar segment 5 enlarged and metasomal sternum 2 with a distinct process. My stuff from Guana and the St. Thomas wasps agree with *spinosa* rather than with *discisa*.

Bembix americana (Fabricius). A female from Anegada I. was identified by Bohart as *B. multipicta* F. Smith; it is the same as my specimens from Guana. Both sexes of the Guana population will go straight to *B. americana* in the key by Evans & Matthews (1968, North American *Bembix*, a revised key and suggested grouping, Annals ESA 61:1284-1299). WHEN WILL SOMEBODY WORK ON CARIBBEAN SPHECIDS AND CLEAR UP SOME OF THESE PROBLEMS?

Got a whole bunch of Bethylidae down there on Guana on these last two trips, mostly from flight traps. A quick glance through this material suggests a minimum of 17 species; my list of Guana aculeates in **Sphecos** 24 included 8 species. I will report further on these cute little kids in a later issue, after I've had a chance to look at them in some detail.

I have made some progress on a puzzling little wasp that superficially looked like a bethylid, but wasn't. It turned out to be a species belonging to the subfamily Amiseginae of the Chrysididae, a group that I've never collected before. So, one more aculeate for Guana Island: Adelphe sp. - 1 female specimen from a Malaise trap in open forest. This may be the first record for the family Chrysididae in this area; I have not yet seen a copy of Kimsey and Bohart (1990. The chrysidid wasps of the world. Oxford Univ. Press. 652 p.). One species of Adelphe has been reported from Cuba. These curious little wasps are egg parasites of phasmatids. I'll know more about this one after I send it to Bohart.

#### The California Deserts by Roy Snelling

Spring in the California deserts was mostly disappointing. Because of the heavy rains last winter, everybody assumed that there would be a great display of flowers and a great collecting season. Well, the great display of flowers generally didn't happen (except in the usual places like Anza-Borrego and Joshua Tree), and the collecting was definitely hard work. My son Gordon and I went out a few times on day-trips; got some pretty good stuff, but had to work hard for all of it. Our best collecting turned out to be on a species of prostrate euphorb (Chamaesyce, species still undetermined): we'd sit down on the ground and aspirate critters coming to these tiny blossoms. An amazing array of species! Karl Krombein (1961, Ent. News 72:80-83) wrote a "quickie" on the results of his casual collecting on C. albomarqinata near the Southwestern Research Station, Portal, Arizona; he recorded 31 species of aculeates, mostly Sphecidae. Gordon and I have

collected about half a dozen times at one spot, near Adelanto in San Bernardino County, dates ranging from 22 May to 14 August. We have a minimum of 75 species. I have only begun to sort the stuff, so cannot now provide anything looking like a list, but can state that the diversity of sphecids is pretty impressive. Most of these species are in the small to tiny range in all groups. In May we picked up many specimens of both sexes of a very small Pseudomethoca. The females were very timid and tended to hide under the euphorb mats when disturbed. The species appears to be undescribed. One female of yet another species was collected at a different site. I expect that Gordon and I will continue to work this resource, probably picking several sites in both the Mojave and Colorado deserts here in California and see what we can come up with. Hey. Wasbauer: got some neat little pepsines, too.

So much for 1993. A very interesting year! I can hardly wait for what 1994 will bring. Stay tuned.

### Mona Island, Puerto Rico - 1992 by Susan C. Jones

USDA Forest Service, Carl Hayden Bee Laboratory, 2000 E. Allen Road, Tucson, AZ 85719

My friend and colleague, Juan A. Torres, just sent me a copy of Roy Snelling's article in the February 1993 Sphecos 24:17-19, which describes several of Roy's collecting trips to the Caribbean. I appreciate the effort that Juan made to arrange an excellent collecting trip to Mona Island and I thank him for inviting me to join his and Roy's expedition during November 1992. I didn't understand the extensive background preparations that Juan had endured, until I tried to arrange another collecting trip to Mona Island in March 1993 (eventually very successful). But that's a story for the International Isoptera Society newsletter.

I must agree with Roy that Mona Island is a wonderful site, both for collecting Hymenoptera, as well as Isoptera, although I find the latter much more interesting. Located 74 km off the west coast of Puerto Rico, Mona Island is a 55 km<sup>2</sup> limestone plateau whose vegetation is categorized within the subtropical dry forest life zone. Its occupants include the giant Mona rock iguana, the Mona boa, four bat species, and introduced feral cats, goats, pigs, mice and rats. Since Roy described our live rat encounter in some detail (and which makes me sound like a wimp), I thought it appropriate to include:

#### "And Now for the Rest of the Tale... El Grande Rata de Isla Mona" (The Big Rat of Mona Island)

I awoke on the fourth morning of our collecting trip to find Juan shaving in the bathroom of the casita that he, Roy, and I shared. There was a full day of collecting ahead, so I decided to make a brief excursion to the mess hall to start my coffee before cleaning up. Had I drank a cup or two of my delicious Louisiana coffee, my suspicions immediately would have been alerted as I reentered our casita, since Roy and Juan hastily exited without our usual prolonged conversation.

Imagine my horror when I walked into the just vacated bathroom and immediately discovered a "dead" rat (at least 8 inches long, excluding the tail) floating in the toilet bowl. I ran outside and called to Roy and Juan that "THERE'S A RAT IN THE TOILET!" Please note that I had yet to be informed of the Gatorade® hypothesis that was formulated (concocted) by my distinguished colleagues. With his usual dry sense of humor, Roy informed me that I was a biologist and should "Deal with it!" Well, I needed to use the bathroom, so I saw little use in a prolonged argument with Roy. I had been taught that the first person to encounter a problem should try to deal with it, rather than leaving it for someone else. Unfortunately for me, some people apparently disagree with this philosophy.

I returned to the bathroom and pondered my dilemma. Should I remove the rat and place it in either Juan's bed or Roy's bed? No. As a biologist, I reasoned that a water burial of Roy's cousin would be the most expedient ... so I flushed ... and the large "dead" rat began scrambling up the sides of the toilet bowl!!!!! I rushed into the outer room and grabbed Roy's walking stick to prod the rat, which eventually disappeared out of sight. I had to repeatedly flush the toilet in order to unplug the pipe. I then thought back to a few hours earlier, when I had trekked to the bathroom in the dark while the generator for electricity was turned off. I'm thankful that the rat hadn't yet taken the plunge into

the bowl, for I would have wakened everyone at Playa Sardinera if that rat had made a move!

After I calmed down, I returned to the mess hall to find my colleagues enjoying a nice breakfast. I was rather upset that, as a third party to the crime, I had to "deal with" the rat. Their rationale - I had left a package of Gatorade®, which the rat had ravaged and then consumed; the rat subsequently had fallen into the toilet trying to quench its powerful thirst. Now, I'm a reasonable person, but this hypothesis requires a considerable stretch of one's imagination. Juan later sent a letter outlining another insightful hypothesis, again based on the Gatorade<sup>®</sup>. He figures that since the rat lived on Mona Island, it couldn't read English and had interpreted the wording as "Gato" (cat, in Spanish), Hence, the vicious attack on the package.

The rest of the tale ... Feral rats and pigs have previously been reported on Mona Island, but I've discovered that this island paradise is frequented by more than one of either species. The reader will have to decide if the title of my story pertains to a 2-legged or 4-legged mammal!

Yes, Roy, I did see some humor in the situation. For Christmas, Roy and Juan each received a plastic rat in the mail.

#### More About Biological Control of Rats by Justin Schmidt Southwestern Biological Institute, 1961 W. Brichta, Tucson, AZ 85745

We aculeatists who fancy wasps should broaden our interests and occasionally come to the aid of humanity as a whole. After all, we owe society a debt for allowing us to indulge in the wonderful activity of studying the most interesting insects on earth (we could be studying whiteflies).

This brings me to the subject of biological control of rats as described by Roy Snelling in Sphecos 24:18. He relates the story on Mona Island of sharing a cabana with Susan Jones (a termitologist), who fed a fine rat specimen her dried Gatorade<sup>®</sup> (for those not familiar with Gatorade<sup>®</sup>, it is a mixture of salts and glucose designed to provide energy and replace the salts lost by sweating athletes). Gatorade<sup>®</sup> is generally consumed with copious quantities of water, but in this case, Susan neglected to inform the rat of that. Thus, after consuming most of the packet of Gatorade® the poor rat probably developed a nasty tummy ache (as the Gatorade acted like a sponge in its stomach), and discovered that Susan had also neglected to provide a water dish. So the poor animal sought the only water available - that in the privy. Unfortunately, the toilet bowl was not shaped with rats in mind, so it had to jump (fall?) in to get a drink. This was its fatal undoing, as it now could not get out. After some fanfare which I will not repeat here, Susan unceremoniously dispatched the rat.

I relate this story because I believe Roy was entirely out of order in picking on poor Susan. Did he not realize the brilliant discovery Susan had made? Did he not realize that she possibly had just made one of the most monumentous leaps forward in our ongoing battle against rats? Rather than making life difficult for her, he should have praised her and proclaimed what a wonderful discovery she had made - a nontoxic and effective means of biological control of rats! Just think what a service Susan has done for humanity to discover this biological control method and possible rid us of rats. Roy, shame on you.

#### "The Gatorade<sup>®</sup> Rat" <u>a Response</u> by Roy Snelling

Dept. of Entomology, Natural History Museum of Los Angeles County, 900 Exposition Blvd., Los Angeles, CA 90007

Here's a comment on the tale of the "Gatorade® Rat" (not related to the Stainless Steel Rat). Susan Jones clearly did not like my account of the incident and put forth her own version. Sheer calumny (how I've awaited an opportunity to use that wonderful word; nobody ever gave me a chance before, inasmuch as everything people have said about me in the past has always been plain, unvarnished truth [and if you believe that maybe we could get together and talk about some real estate deals I have in mind]). Arnold very kindly sent me an advance copy, thereby permitting me to respond to this dastardly and unwarrented attack upon both my character and veracity.

Susan's telling does twist the tale/ tail. Imprimis: When Susan prepared to enter the Place of the Rat, Juan and I did not immediately exit the cabaña (= "casita" of her version), inasmuch as I was already without and Juan, in fact, had earlier repaired to the cocina for a first cup of coffee. I was on my way there when she yelled at me to come and get rid of that "big rat in the toilet." Secundus: I certainly did not imply that she was a "wimp". Indeed, quite the opposite: I informed her that as a biologist she should be able to take care of the problem, thus making clear my full faith in her machisma. As for the rat being a problem: I encountered no difficulty whatsoever with respect to the small mammal (and it truly was not a very large rat), nor did Juan (and she failed to comment to the fact that it was Juan who saw it first, not I). She is the one who for some reason I comprehend found it to be problematic. Tertius: what "walking stick"? Neither Juan nor I (both older than she) uses a "walking stick". What rot! Susan may think she's a reasonable person - maybe she is. However, before casting unfounded and unsupported aspersions on the Gatorade® hypothesis, I suggest she consume an ounce or two, sans H<sub>2</sub>O; she might well then be inclined to ingest a drop or two of that sublime liquid that led to the downfall of our not-much-lamented cousin. On the other hand, that was a darksome and cruel deed she did; the poor water-soaked rodent did her no harm, yet she pitilessly put it to death by drowning, rather than solicitously lifting it from its dire peril by means of that handy handle, the tail. This heartless person even took the trouble to determine that the inoffensive animal was female (rata); probably she orphaned a whole litter of dependant youngsters, too! Other, rather minor, imperfections in her account of "El Grande Rata de Isla Mona" (Juan, I'll bet you didn't believe that she knew Spanish) I shall let pass. But, what can you expect from someone who thinks that those pale, squishy termites are more interesting than ants and other Hymenoptera?

### MUSEUM/COLLECTION NEWS

Types In the Manfredo Fritz Collection by Manfredo Fritz Casilla Correo 539 4400 Salta, Argentina

The following is a list of the types in my collection, comprising 137 Holotypes and 30 Allotypes. I intend to donate all these types to the Museo Argentino de Ciencias Naturales in Buenos Aires in the near future. As you know, two young, enthusiastic hymenopterists [Genise and Roig] are working intensively in that institution with wasps and bees, and I think this is the best heritage that I can leave for them.

#### PERGIDAE

Lagideus ucratus D.R. Smith – Holotipo 1990. Revta bras. Ent. 34:123.

#### CHALCIDIDAE

Leucospis leptomera Boucek - Holotipo 1976. Studia Ent. 17:430.

### **ICHNEUMONOIDEA**

- Stephanus alvarengai Orfila Holotipo 1960. Actas Ier. Congr. Sudam. Zool. 3:125.
- Stephanus brasiliensis Orfila Holotipo 1960. Actas Ier. Congr. Sudam. Zool. 3:132.

*e orfilai* nom.n. De Santis, 1980. Cat. Hym. Bras. Parasitica p. 8 (1982)

Stephanus camposseabrai Orfila - Holotipo

1959. Rev. Soc. Ent. Arg. 21:39.

Stephanus carioca Orfila – Holotipo 1960. Actas ler. Congr. Sudam. Zool. 3:129.

Stephanus sanmartinianus Orfila - Holotipo

1949. Acta Zool. Lilloana 7:350 Stephanus willineri Orfila – Holotipo

1949. Acta Zool. Lilloana 7:345 Neosteohanus fritzi Orfila - Holotipo

Neostephanus fritzi Orfila – Holotipo 1956. Rev. Soc. Ent. Arg. 19:6.

### CHRYSIDIDAE

Cleptes (Neocleptes) fritzi Kimsey – Holotipo

1981. Proc. Biol. Soc. Wash. 94:816. *Ipsiura fritzi* Bohart – Holotipo 1985. J. Kansas Ent. Soc. 58:715.

#### TIPHIIDAE

Anthosila argentina Genise – Holotipo 1984. Rev. Soc. Ent. Arg. 43:216. *Tiphiodes chaco* Genise – Holotipo 1984. Rev. Soc. Ent. Arg. 43:209. *Tiphiodes toba* Genise – Holotipo 1984. Rev. Soc. Ent. Arg. 43:216.

#### ORUSSIDAE

Guiglia chilensis Benson – Alotipo 1955. Proc. r. ent. Soc. Lond. B 24:112.

### BRADYNOBAENIDAE

Bradynobaenus bidentatus Genise 1986. Physis 44(C):50. – Holotipo Bradynobaenus cordobensis Genise 1986. Physis 44(C):52. – Holotipo

### MUTILLIDAE

Eotilla typhoctoides Martínez & Fritz 1975. Rev. Chil. Ent. 8:23. - Holotipo Euspinolia casali Martínez & Fritz 1976. Neotrópica 19:113. - Holotipo Lophomutilla calva Fritz - Holotipo 1990. Rev. Soc. Ent. Arg. 48:130. Lophomutilla ophomuti Fritz - Holotipo 1990. Rev. Soc. Ent. Arg. 48:132. Lophomutilla suarezi Fritz & Pagliano - Holotipo 1993. Boll. Soc. ent. ital. 124:214. Lophomutilla braulioi Fritz & Pagliano - Holotipo 1993. Boll. Soc. ent. ital. 124:213. Lophomutilla inca Fritz & Pagliano Holotipo 1993. Boll. Soc. ent. ital. 124:215. Lophomutilla vina Fritz & Pagliano Holotipo 1993. Boll. Soc. ent. ital. 124:213. Lophomutilla correntina Fritz & Pagliano - Holotipo 1993, Boll, Soc. ent. ital, 124:212, Lophomutilla oscura Fritz & Pagliano - Holotipo 1993. Boll. Soc. ent. ital. 124:213. Pertyella aguaz Fritz Holotipo 1990. Rev. Soc. Ent. Arg. 48:134. Pertvella cordoi Fritz - Holotipo 1990. Rev. Soc. Ent. Arg. 48:135. Pertyella diasi Fritz Holotipo 1990. Rev. Soc. Ent. Arg. 48:137. Pertyella uzai Fritz Holotipo 1990. Rev. Soc. Ent. Arg. 48:137. Pertyella guarani Fritz & Pagliano Holotipo 1993. Boll. Soc. ent. ital. 124:218. Lvnchiatilla tacana Casal Alotipo

1963. Acta Zool. Lilloana 19:365.

### EUMENIDAE

Discoelius schajovskoyi Willink - Alotipo 1956. Neotrópica 2:55. Montezumia fritzi Willink

- Holotipo, Alotipo 1982. Bol. Ac. Nac. Cienc. Córdoba 55:164.

### SPHECOIDEA -- LARRINAE

Pisonopsis australis Fritz – Holotipo, Alotipo 1968. Neotrópica 11:53. Tachytes fritzi Bohart – Holotipo 1979. Tr.Am.Ent.Soc. 104:464. Tachytes rufalaris Bohart – Holotipo 1979. Tr. Am. Ent. Soc. 104:484. Tachytes willinki Bohart – Holotipo 1979. Tr. Am. Ent. Soc. 104:500.

#### SPHECOIDEA - PEMPHREDONIDAE

Pluto colonensis van Lith	- Holotipo
1979. Tijd. Ent. 122:198.	
Pluto fritzi Van Lith	- Holotipo
1979. Tijd. Ent. 122:199.	
Pluto obscurus Van Lith	– Holotipo
1979. Tijd. Ent. 122:214.	
Psen (Pseneo) nigripes var	n Lith
1975. Ent. Ber. 35:171.	- Holotipo
Psen (Pseneo) auriger Van	Lith
- Hold	tipo, Alotipo
1975, Tiid, Ent. 118:25,	-

#### SPHECOIDEA - SPHECINAE

- Isodontia alvarengai Fritz Holotipo 1983. Rev. Soc. Ent. Arg. 42:28. Isodontia guaranitica Willink – Holotipo
- 1951. Acta zool. lilloana 11:92.

Isodontia paranensis Berland – Alotipo 1951. Willink. Acta zool. Lilloana 11:85. [this is not a true allotype - editor]

#### SPHECOIDEA - NYSSONINAE

Antomartinezius boharti Fritz – Holotipo 1974. An. Mus. Hist. Nat. Valparaíso 6:198. Foxia deserticola Fritz – Holotipo

- 1959. Rev. Soc. Ent. Arg. 21:133. Foxia garciai Fritz – Holotipo
- 1972. Acta Scient. (Ent.) 6:9. Foxia martinezi Fritz - Holotig
- Foxia martinezi Fritz Holotipo 1972. Acta Scient. (Ent.) 6:7. Foxia pirita Fritz – Holotipo
- Foxia pirita Fritz Holotipo 1972. Acta Scient.(Ent.) 6:6. Foxia tercera Fritz – Holotipo
- 1972. Acta Scient.(Ent.) 6:4. Idionysson cordialis Fritz – Holotipo
- 1970. An. Mus. Hist. Nat. Valparaíso 1:150.

- Losada penai Fritz Holotipo 1974. An. Mus. Hist. Nat. Valparaíso 6:200.
- Metanysson carcavalloi Fritz Holotipo 1959. Rev. Soc. Ent. Arg. 21:134.
- Metanysson cicheroi Fritz Holotipo 1974. An. Mus. Hist. Nat. Valparaíso 6:194.
- Metanysson diezguitas Fritz Holotipo 1956. Neotrópica 2:94.
- Metanysson foersteri Fritz Holotipo 1956. Neotrópica 2:92.
- Metanysson fratemus Fritz Holotipo 1970. An. Mus. Hist. Nat. Valparaíso 1:144.
- Metanysson horacioi Fritz Holotipo 1974. An. Mus. Hist. Nat. Valparaíso 6:191.
- Metanysson toba Fritz Holotipo 1974. An. Mus. Hist. Nat. Valparaíso 6:195.
- Metanysson tropicalis Fritz Holotipo 1970. An. Mus. Hist. Nat. Valparaíso 1:146.
- Nysson (Epinysson) casali Fritz
- Holotipo, Alotipo 1970. An. Mus. Hist. Nat .Valparaíso 1:148.
- Heliocausus latitarsus Fritz & Toro – Holotipo, Alotipo
- 1977. Rev. Soc. Ent. Arg. 35:26. Gorytes bruchi Schrottky – Holotipo
- Fritz. 1964. Neotrópica 10(33):108. Gorytes cordobensis Fritz

- Holotipo, Alotipo

- 1964. Neotrópica 10:110.
- Gorytes schrottkyi Fritz Holotipo, Alotipo 1964. Neotrópica 10:109.
- Harpactostigma Ilanoi Fritz
  - Holotipo, Alotipo 1964. Neotrópica 10:54.
- Harpactostigma patagonica Fritz
- Holotipo, Alotipo 1959. Rev.Soc.Ent.Arg. 21:135.
- Bembecinus comechingon Willink

- Holotipo

- 1949. Acta zool.lilloana 7:108. Bembecinus bridarollii Willink – Holotipo
- 1949. Acta zool. lilloana 7:106. Bembix latigenata Willink – Holotipo 1976. Acta zool. lilloana 4:590.

#### SPHECOIDEA - CRABRONINAE

- Anacrabro fritzi Leclercq Holotipo 1973. Acta zool. lilloana 30:46.
- *Ectemnius servitorius* Leclercq Holotipo 1972. Bull. Soc. R. Sc. Liege 41: 203-218.
- Lecrenierus piraponensis Leclercq
  - Holotipo 1979. Bull. Rech. Agron. Gembloux 9:67.

Podagritus bordai Fritz - Holotipo, Alotipo 1971, Rev. Soc. Ent. Arg. 33:112. Podagritus carrascoi Fritz - Holotipo, Alotipo 1971. Rev. Soc. Ent. Arg. 33:110. Podagritus valenciai Fritz - Holotipo, Alotipo 1971.Rev. Soc. Ent. Arg. 33:113. Podagritus (Chilichuca) brieni Leclercq - Holotipo 1981, Ann. Soc. r. Zool. Belg. 111:76. Podagritus (Parechuca) quiacae - Holotipo Leciercq 1982. Ann. Soc. r. Zooi. Belg. 112:274. SPHECOIDEA - PHILANTHINAE Odontosphex fritzi Menke - Holotipo 1967. Pan-Pac. Entom. 43:145. Cerceris abac Fritz & Mariluis - Holotipo 1977. Rev. Soc. Ent. Arg. 35:100. Cerceris and algalensis Fritz & Toro - Holotipo 1969. An. Mus. Hist. Nat. Valparaíso 2:153. Cerceris apix Fritz & Mariluis -Holotipo 1979. Acta Scient. (Ent.) 12:34. Cerceris brethesi Fritz & Mariluis - Holotipo 1979. Acta Scient. (Ent.) 12:34. Cerceris caiman Fritz & Mariluis - Holotipo 1977. Rev. Soc. Ent. Arg. 35:100. Cerceris carrascoi Fritz & Toro - Holotipo, Alotipo 1974. An. Mus. Hist. Nat. Valparaíso 6:182. Cerceris caura Fritz & Mariluis - Holotipo 1977. Rev. Soc. Ent. Arg. 35:111. Cerceris cuica Fritz & Mariluis - Holotipo, Alotipo 1977, Rev. Soc. Ent. Arg. 35:109 Cerceris curita Fritz & Mariluis - Holotipo, Alotipo 1979. Acta Scient. (Ent.) 12:30. Cerceris dione Fritz - Holotipo 1961. Actas Trab. ler. Congr. Sudam. Zool. 3:67. Cerceris lamarquensis Fritz & Toro - Holotipo 1969, An. Mus. Hist, Nat. Valparaíso 2:142. Cerceris manca manca Fritz & Mariluis - Holotipo 1979. Acta Scient. (Ent.) 12:33. Cerceris manca cordil Fritz & Mariluis - Holotipo 1979. Acta Scient. (Ent.) 12:33. Cerceris liubae Fritz - Holotipo 1990. Rev. Soc. Ent. Arg. 48:154. Cerceris manjoni Fritz & Mariluis - Holotipo, Alotipo

1977. Rev. Soc. Ent. Arg. 35:99. Cerceris mariusi Fritz –\_Holotipo 1989. Insecta Mundi 3(4):262.

Cerceris menkei Fritz & Toro – Holotipo 1969. An. Mus. Hist. Nat. Valparaíso 2:151.

Cerceris narcisoi Fritz – Holotipo 1983. Rev. Soc. Ent. Arg. 42:164.

Cerceris nigrosa Fritz – Holotipo 1989. Insecta Mundi 3(4):263.

Cerceris paulista Fritz – Holotipo 1983. Rev. Soc. Ent. Arg. 42:168.

Cerceris penai Fritz & Toro – Holotipo 1969. An. Mus. Hist. Nat. Valparaíso 2:143.

Cerceris sanluis Fritz & Toro – Holotipo 1969. An. Mus. Hist. Nat. Valparaíso 2:147.

Cerceris serrana Fritz – Holotipo 1989. Insecta Mundi 3(4):265.

Cerceris tingi Fritz & Mariluis –Holotipo 1977. Rev. Soc. Ent. Arg. 35:104.

Cerceris vianai Fritz & Toro – Holotipo 1969. An. Mus. Hist. Nat. Valparaíso 2:153.

Cerceris willineri Fritz – Holotipo 1961. Act. Trab. ler. Congr. Sudam. Zool. 3:70.

Cerceris willinki Fritz – Holotipo 1983. Rev. Soc. Ent. Arg. 42:164. Cerceris zain Fritz – Holotipo

1990. Rev. Soc. Ent. Arg. 48:156.

Cerceris zelichi Fritz & Mariluis - Holotipo, Alotipo

1977, Rev. Soc. Ent. Arg. 35:106.

#### APOIDEA

Coelioxys genisei Toro & Fritz - Holotipo, Alotipo 1991. Acta Ent. Chil. 16:74. Coelioxys ljuba Toro & Fritz - Holotipo 1991. Acta Ent Chil. 16:75. Coelioxys palmaris Fritz & Toro - Holotipo 1990. Gayana 54(1-2):18. Coelioxys pomona Toro & Fritz - Holotipo 1991. Acta Ent. Chil. 16:78. Coelioxys roigi Fritz & Toro - Holotipo 1990. Gayana 54(1-2):16. Coelioxys ruzi Toro & Fritz Holotipo, Alotipo 1991, Acta Ent. Chil. 16:78 **Dialictus flavipes Moure** Holotipo 1950. Dusenia 1:307. Ceratalictus allostictus Moure - Holotipo, Alotipo 1950. Dusenia 1:318. Oxaea stenocoryphe Moure - Holotipo, Alotipo 1947. Rev. Soc. Ent. Arg. 13:223. Leptometria (Leptorachis) bridarollii Moure - Holotipo

1947. Rev. Soc. Ent. Arg. 13:237. Leptometria willineri Moure - Holotipo, Alotipo 1947. Rev. Soc. Ent. Arg. 13:239. SPECIES IN PRESS OR WITHOUT DATE OF PUBLICATION - Holotipo Apterodryinus fritzi Olmi Lynchiatilla tuzai Fritz Holotipo - Holotipo Horcomutilla ypane Fritz Horcomutilla willineri Fritz - Holotipo Xystromutilla tingoensis Fritz - Holotipo Megischus townesi Fritz & Scaramozzino - Holotipo Hemistephanus amazonicus Fritz & - Holotipo Scaramozzino Hemistephanus bolivianus Fritz & Holotipo Scaramozzino Hemistephanus cariocanus Fritz & Scaramozzino Holotipo Hemistephanus orfilanus Fritz & Scaramozzino - Holotipo Coelioxys quechua Toro & Fritz - Holotipo Coelioxys balastos Toro & Fritz - Holotipo Coelioxys trancas Toro & Fritz - Holotipo, Alotipo Coelioxys noa Toro & Fritz - Holotipo, Alotipo Coelioxys alisal Toro & Fritz - Holotipo Coelioxys saltensis Toro & Fritz - Holotipo Coelioxys tastil Toro & Fritz - Holotipo - Holotipo Trimeria bequaerti Willink Paragapostemon implexus Moure - Holotipo Psaenythia capnistoptera Moure - Holotipo

1947. Rev. Soc. Ent. Arg. 13:234.

Dichanthidium exile Moure

Crossocerus annulipes (Lepeletier & Brullé), female (Sphecidae), North America.

- Holotipo, Alotipo

# **NEW BOOKS**

Hymenoptera of the World. An Identification guide to families. Edited by Henry Goulet and John T. Huber. Agriculture Canada Publication 1894/E. Ottawa, Canada, 1993. 668 pages, many illustrations. \$63.35 in US dollars (for orders outside of Canada), or \$48.75 in Canadian dollars (for Canadian orders) (see Sphecos 25:22 for ordering information).

This impressive tome with its impressive title has finally appeared. Hymenoptera of the World is printed in large format, 8 1/2 by 11 inches, 99 families of Hymenoptera are recognized by the 11 authors, not all of whom are Canadian, but all of whom are experts in their respective sections. The book is aimed at the novice user (non-taxonomists and students), and therefore, the identification keys are illustrated lavishly with figures. The keys go to the level of subfamily for aculeates and ichneumonoids. Family by family treatments follow the keys and these include habitus figures. diagnoses, statistics on included taxa. distribution, biological notes and important references (mostly taxonomic). The numerous habitus figures are printed one per page resulting in a lot of wasted page space; the figures are so large that Hymenoptera of the World is already being referred to by some hymenopterists as the "coloring book". These drawings could have been reduced in size and grouped several to a page without loss of clarity. This would have reduced the number of pages of the book and reduced production costs.

Introductory chapters present a good overview of the order Hymenoptera (which includes a list of newsletters and their editors, etc., a nice touch), and a survey of morphology. The latter is very well illustrated so that anyone should be able to grasp terminology. In fact the chapter ends with a 25 page illustrated Glossary in which nearly every term is defined and shown by an accompanying figure. Excellent!

In Hymenoptera there are often a number of different terms for the same structure, as the authors of the morphology chapter point out. An attempt has been made here to standardize morphological terms using "the simplest acceptable terms", a highly laudible goal. But the aculeate section of the book, a significant hunk of Hymenoptera of the

World, does not adhere strictly to this goal. For example, the system for naming wing cells in aculeates does not follow the Comstock/Needham system employed in the Glossary. Instead cell names follow those in The Hymenoptera by Gauld and Bolton, 1988 (which is similar to the names used in Sphecid Wasps of the World by Bohart and Menke, 1976, and other aculeate literature). As usual, aculeate workers lead the way. We were glad to see that standardization resulted in universal usage of pedicel, scape and flagellomeres in this book. No more anelli, ring segments, and so on in Chalcidoidea. Nevertheless the Glossary contains two different terms for one mesopleural area: prepectus and epicnemium. The epicnemial carina of the mesopleuron was called the omaulus in Sphecid Wasps of the World. We wonder why omaulus was not used in the Glossary when sternaulus was accepted there (probably because it is also used in some Ichneumonoidea). Use of the word torulus for the antennal socket seems unnecessary and less descriptive to us, and in fact the authors themselves use socket instead of torulus in explaining the term radicle. Not surprisingly the Glossary espouses mesosoma and metasoma instead of thorax and abdomen. The more descriptive term plantar lobe is used in place of plantula. Following the fairly recent work of Wootton (1978) on wings, the anal lobe of the hindwing is called the claval lobe in the Glossary. This reflects the fact that, in general, the morphology section is quite up-to-date.

The keys are certainly user-friendly. Couplet characters are listed line by line, each preceded by a letter. Figures illustrating these characters and indexed by the couplet number and character letter, are inserted above their respective couplet. Thus the user can quickly and easily see a particular character. This has resulted in a novel type of key construction, and it works very well. The figures are mislabeled on page 302. The scutum should be "b" and "bb", and the lateral views of the head and thorax should be "c" and "cc".

The organization of superfamilies departs from tradition. The "sawfly" groups come first, followed by the Aculeata and then the Parasitica. In the discussion of the monotypic living fossil Anaxyelidae no mention is made of the other name sometimes improperly used for this family, Syntexidae. Literature references in

the various family treatments include papers published as recently as 1993, but we noted significant omissions. Missing from the tiphiid references was the 1992 Reclassification of Tiphiidae with description of a new subfamily from Turkey by Argaman and Özbek, and Genise's 1992 Las especies conocidas de Tiphiidae de la Argentina y paises vecinos. Under Sierolomorphidae no mention is made of Argaman's 1991 paper Generic synopsis of Sierolomorphidae in which he recognized two genera. California spider wasps of the subfamily Pompilinae. 1985, by Wasbauer and Kimsey was not cited under Pompilidae, nor were any of Heinrich Wolf's significant papers on this family.

Recognition of one superfamily, Apoidea, for sphecid wasps and bees follows contemporary thinking and is nice to see here. Following the lead of Brothers' now classic 1975 aculeate phylogeny paper, the Apoidea is divided into two informal groups, Spheciformes for sphecid wasps, and Apiformes for bees. But if only two families are recognized, Sphecidae and Apidae, these suprafamilial names would be unnecessary. Albert Finnamore, author of the Spheciformes section, elevates the subfamilies recognized in Sphecid Wasps of the World to family (with minor exceptions), stating that the rationale for doing so is to "make the classification comparable to that widely accepted in the Apiformes". We wonder why this argument is never reversed: why not make bees comparable to sphecids and recognize only Apidae (as is done by Gauld and Bolton in The Hymenoptera, and by other European workers). Finnamore says that none of the current higher category systems for Sphecidae s.l. are likely to be correct: thus his elevation of sphecid subfamilies to families seems premature and particularly inappropriate in an identification manual.

In the sphecid section, the group with which we are most familiar, we noted a number of shortcomings. The key presented on page 279 that separates Spheciformes from Apiformes uses traditional characters and omits one of the best, namely the presence in sphecids (and absence in bees) of a cleaning pecten or brush on hindtarsomere I and its apposing pectinate tibial spur. Finnamore has made a number of unilateral higher level changes in Spheciformes without discussing why. It seems to us that such changes are inappropriate in a book whose main purpose is identification rather than phylogeny. The Laphyragoginae has been included in his Astatidae, and the Xenosphecinae was put in Mellinidae without explanation. Furthermore, there is no discussion of recognizing Mellinidae as an entity separate from Nyssonidae. On the other hand, Finnamore places Entomosericinae in Nyssonidae and his rationale for doing so is explained. Larrinae and Crabroninae were combined as one family, Crabronidae, and although unification of these two groups has been proposed by others (Evans, 1964, Lomholdt, 1985 and Menke, 1988), no mention is made of these recent papers. Furthermore, usage of the family group name Crabronidae for this assemblage, instead of Larridae, was not explained. As it turns out, Crabronidae is the older of the two names. Why was this not discussed?

Apparently species totals for families of Spheciformes were taken from Sphecid Wasps of the World without adding taxa described since that book was published nearly 18 years ago. Thus the totals in most cases are out-of-date. For example, 195 species are currently known in the Ampulicinae, 1435 in the Crabroninae and 2572 in the Larrinae (rather than 167, 1300 and 2000, respectively.) Pulawski (1978) was the author of the European USSR paper cited on page 280, not Tobias. Generally the literature for Spheciformes is up-to-date but the following should have been included: Dollfuss' 1991 Bestimmungsschlussel der Grabwespen Nord- un Zentraleuropas. Under Heterogynaidae Argaman's 1986 paper Taxonomy of Heterogynaidae was not cited and it included one new genus and a key to genera. Also the group is known from Turkmenistan as indicated by Antropov and Gorbatovsky (1992). Finnamore states that "no major works have appeared for Sphecidae [i.e., Sphecinae] since Bohart and Menke (1976)" but this is simply not true. For example Vardy revised Trigonopsis in 1978, and Hensen revised the subgenus Prosceliphron in 1987, and the genus Chalybion in 1988. Literature for Pemphredonidae is very incomplete; only one paper by Finnamore is cited. Menke (1989) established a new subtribal classification for part of the Pemphredoninae, and van Lith (1979) and Finnamore (1983) both published keys to species of some genera of Pseninae. Eighme (1989) revised the North American Diodontus, Vincent (1979) revised

the North American Passaloecus, among other papers that could be mentioned for the Pemphredoninae. Matthews' (1968, etc.) significant papers on sociality of Microstigmus are not cited either. Something about the biology of Laphyragoginae is known contrary to what is said on page 296. Kazenas (1985) discovered that one species preved upon moths of the genus Crambus. The improper spelling Trypoxylonini is used on pages 296-297 instead of the correct Trypoxylini. References for Crabronidae omits significant revisions and identification keys such as, for example, Bohart (1976), Crabro, (1979), Tachytes, Coville (1982), Trypoxylon, Krombein and Gingras (1984), Liris, Pulawski (1988). Tachysphex, Menke (1988), Pison, (1992), Larra, etc., etc. To the references for Nyssonidae should be added papers by Bohart (1984), Austrogorytes and Stubblefield (1984), Stizus. Lastly, Rubio's (1976) revision of Trachypus was omitted from the pertinant literature for Philanthidae, as were papers on Cerceris by Evans, Fritz, Tsuneki and others. Eremiaspheciini was misspelled in the key and in the sketch.

Hymenoptera of the World is a substantial work, and should be very useful to the audience for which it was designed. However, it will also be a handy reference for hymenopterists everywhere. Nowhere else will you find well illustrated, modern identification keys to all of the families of Hymenoptera. If you are a hymenopterist, it should be on your bookshelf.

#### A. S. Menke & W. J. Pulawski

#### Literature cited

- Antropov, A. V. and V. V. Gorbatovsky, 1992. *Heterogyna polita* sp. n. (Hymenoptera, Sphecidae) – First representative of the subfamily Heterogynainae in the fauna of the USSR. Bull. Moscov. obshch. ispyt. prirody Otd. Biol. 97:53-59.
- Argaman, Q., 1986. Taxonomy of Heterogynaidae. Israel J. Ent. 14:7-12.
- Argaman, Q., 1991. Generic synopsis of Sierolomorphidae. Israel J. Ent. 24: 29-33.
- Argaman, Q. and H. Özbek, 1992. Reclassification of Tiphiidae with description of a new subfamily from Turkey. Türk. Ent. Derg. 16:3-12.
- Bohart, R. M., 1976. A review of the nearctic species of Crabro. Trans.

Amer. Ent. Soc. 102:229-287.

- Bohart, R. M., 1979. *Tachytes* of South America. Trans. Amer. Ent. Soc. 104: 435-505.
- Bohart, R. M., 1984. A revision of the genus *Austrogorytes* Bohart. Aust. J. Zool. 32:391-412.
- Bohart, R. M. and A. S. Menke, 1976. Sphecid wasps of the world, a generic revision. Univ. of California Press, Berkeley. 695 p.
- Brothers, D., 1975. Phylogeny and classification of the aculeate Hymenoptera, with special reference to Mutillidae. Univ. Kansas Sci. Bull. 50:483-648.
- Coville, R., 1982. Wasps of the genus *Trypoxylon* subgenus *Trypargilum* in North America. Univ. Calif. Pub. Ent. 97:1-147.
- Dollfuss, H., 1991. Bestimmungsschlüssel der Grabwespen Nord- und Zentraleuropas (Hymenoptera: Sphecidae). Stapfia, no. 24, 247 p.
- Eighme, L., 1989. Revision of *Diodontus* in America north of Mexico. Ann. Ent. Soc. Amer. 82:14-28.
- Evans, Howard E., 1964. The classification and evolution of digger wasps as suggested by larval characters (Hymenoptera: Sphecoidea). Ent. News 65:225-237
- Finnamore, A. T., 1983. Revision of the American species of *Mimesa*. Lyman Ent. Mus. Res. Lab. Mem. 12, 171 p.
- Gauld, I. and B. Bolton, editors, 1985. The Hymenoptera. Oxford Univ. Press, Oxford. 332 p.
- Genise, J. F., 1992. Las especies conocidas de Tiphiidae de la Argentina y países vecinos. Rev. Soc. Ent. Arg. 50:277-329.
- Hensen, R., 1987. Revision of the subgenus *Prosceliphron* van der Vecht. Tijdschrift v. Ent. 129:217-261.
- Hensen, R., 1988. Revision of the nominate subgenus *Chalybion* Dahlbom. Tidschrift v. Ent. 131:13-64.
- Kazenas, V., 1985. The prey of *Laphyragogus turanicus* Gussakovskij. Sphecos 10:17.
- Krombein, K. V. and S. Gingras, 1984. Revision of North American *Liris* Fabricius. Smithsonian Contrib. Zool. 404, 96 p.
- Lomholdt, O., 1985. A reclassification of the larrine tribes with a revision of the Miscophini of southern Africa and Madagascar. Ent. Scand., Suppl. 24, 183 p.
- Matthews, R. W., 1968. Nesting biology of the social wasp *Microstigmus comes*. Psyche 75:23-45.

- Menke, A. S., 1988. *Pison* in the New World: a revison (Hymenoptera: Sphecidae: Trypoxylini). Contrib. Amer. Ent. Inst. 24:1-171.
- Menke, A. S., 1989. *Arpactophilus* reassessed, with three bizarre new species from New Guinea. Invert. Taxon. 2:737-747.
- Menke, A. S., 1992. Mole cricket hunters of the genus *Larra* in the New World. J. Hym. Res. 1:175-234.
- Pulawski, W. J., 1988. Revision of North American *Tachysphex* wasps including Central American and Caribbean Species. Mem. Calif. Acad. Sci. no. 10, 211 p.
- Rubio, E., 1976. Revisión del género *Trachypus* Klug. Rev. Fac. Agron. Univ. Zulia. 3(1):7-87.
- Stubblefield, J. W., 1984. Genus *Stizus* Latreille in North America. Bull. Mus. Comp. Zool. 150:397-428.
- van Lith, J. P., 1979. The New World genus *Pluto*. Tijdschrift v. Ent. 122: 127-239.
- Vardy, C., 1978. A revision of the neotropical wasp genus *Trigonopsis* Perty. Bull. British Mus. (Nat. Hist.), Ent. Ser. 37:117-152.
- Vincent, D, 1979. A revision of the genus *Passaloecus* in American north of Mexico. Wasmann J. Biol. 36:127-198.
- Wasbauer, M. and L. Kimsey, 1985. California spider wasps of the subfamily Pompilinae. Bull. Calif. Insect Surv. 26:1-130.
- Wootton, R. J., 1978. Function, homology and terminology in insect wings. Syst. Ent. 4:81-93.

Hyménoptères Sphecidae d'Europe Occidentale, Volume 1, Generalites – Crabroninae by Jacques Bitsch and Jean Leclercq. Fauna de France, vol. 79, 325 p. (Available from Faune de France, B.P. N° 392, F-75232 Paris Cédex 05, France. Price: France: FF 441 TTC (TVA au taux actuel de 5,5%) + emballage et port; foreign countries: FF 418 + wrapping and expedition.)

Although this book is part of the Fauna de France series, its scope includes western Europe. Thus it is useful to a far larger audience. The book is only the first part of several (?) anticipated to complete a survey of Sphecidae. Part one includes an introduction to the family and a taxonomic treatment of the subfamily Crabroninae. The introductory section includes a fine presentation on morphology and terms, augmented with illustrations. Terminology follows that of *Sphecid Wasps of the World* with minor exceptions. It is nice to see such continuity. There are tables comparing terms used by different authors for wing and genitalic structures. No mention is made of the changes in wing terminology proposed by Wootton (1978), or the new theory concerning the nature of the propodeal dorsum (metapostnotum) by Brothers (1976). The morphology of larvae is described and illustrated.

There is a key to aculeate families found in Europe, and then a section on classification of Sphecidae. The last includes a historic overview and a chart showing the subfamilies and tribes recognized by the authors. This follows the same arrangement found is Sphecid Wasps of the World except that Dinetini is placed under Larrinae. It is not clear if that was intentional or a copy editor's lapsus. The tribe Mellinini is misspelled "Mellini". No mention is made of the contemporary views of Lomholdt (1985) and Menke (1988), both of whom espoused combining Larrinae and Crabroninae as one subfamily, a change that Evans (1964) proposed 30 years ago. Evans' subfamily and tribal arrangement is given by Bitsch and Leclercq, but not discussed. Since Crabroninae comprises the entire taxonomic treatment of this book, it would have been nice to see some discussion of the merits of combining Larrinae and Crabroninae.

A section on phylogeny follows, and Bitsch and Leclercq have given an upto-date historic overview here. Included is a simplified cladogram from Alexander's (1990) preliminary cladistic analysis of Sphecidae and bees in **Sphecos** 20. Alexander's (1992) more thorough analysis is cited by Bitsch and Leclercq, but the paper probably appeared too late to incorporate a cladogram from it.

Nesting behavior of Sphecidae is summarized. Distribution of the family in France and the procedure for mapping species is discussed. Records for each species treated are shown on a map of France that depicts all the departments. It is at that level that species distribution is plotted. The section on distribution also includes a list of French museums and private collections, and a comprehensive list of literature dealing with sphecid distribution in other parts of Europe.

The systematics section occupies most of the book, and it begins with a key to the sphecid genera known in Europe. This key is illustated and it looks like it should work well. Genera not recorded from France are surrounded by brackets. The key is followed by accounts of the tribes Crabronini and Oxybelini and their genera and species. The genera are treated in alphabetical order. For each genus there is a list of synonyms (if any) and important literature. These are followed by a description of morphology, and a summary of known biology with citations of papers. The number of species known for Europe, France and other areas of the world is given, and taxonomic problems are discussed (recognition of subgenera, etc.). A key to subgenera is provided if they are recognized, and, of course, there is a key to species for each genus. The species keys are well illustrated and they should prove very useful throughout western Europe, and possibly elsewhere.

Individual species treatments follow the same format used for genera. Occasionally subspecies are recognized and they are keyed as well. In view of the morphological differences mentioned for some subspecies, I wonder if they aren't really species?

After the literature cited section there is a page giving new synonymy proposed in this book, as well as a several page floral record list by wasp species.

In summary, this is a very fine, up-todate treatment of the Crabroninae of Europe, and a nice introduction to the Sphecidae. The well illustrated identification keys should find a wide appeal in the user community. The authors are to be congratulated on producing such a useful book. I hope that the remaining sphecid groups will get the same treatment soon.

#### Arnold S. Menke

#### Literature cited:

- Alexander, B. A., 1990. A preliminary phylogenetic analysis of sphecid wasps and bees. Sphecos 20:7-16.
- Alexander, B. A., 1992. An exploratory analysis of cladistic relationships with the superfamily Apoidea, with special reference to sphecid wasps. J. Hym. Res. 1:25-61.
- Brothers, D. J., 1976. Modifications of the metapostnotum and origin of the 'propodeal triangle' in Hymenoptera Aculeata. Syst. Ent. 1:177-182.

- Evans, H. E., 1964. The classification and evolution of digger wasps as suggested by larval characters. Ent. News 75:225-237.
- Lomholdt, O., 1985. A reclassification of the larrine tribes with a revision of the Miscophini of southern Africa and Madagascar. Ent. Scand., Suppl. 24, 183 p.
- Menke, A. S., 1988. *Pison* in the New World: a revision. Contrib. Amer. Ent. Inst. 24:1-171.
- Wootton, R. J., 1978. Function, homology and terminology in insect wings. Syst. Ent. 4:81-93.

# PURSUING LEFT-HANDED ENTOMOLOGISTS

When I finished reading the comments on left-handed entomologists (Sphecos 23:15) I thought that this idea could very well be pursued. However, this turned out not to be the case.

In talking with fellow entomologists I found that most are right-handed and do place labels correctly (i.e. they can be read without rotating the pin with the fingers). One of them always places the labels in the inverted position; this person is right-handed, but his dominant eve is that of a left-handed person. I was surprised by the chaos while checking the collection of one of Cuba's best entomologists, Dr. Pastor Alayo. He is left-handed and in his collection labels are oriented in different ways. He told me that his main concern is not the orientation of the text, but rather that the pin does not puncture (and consequently does not deface) the written information.

Not one of the labelers has been guided by an entomological text book.

From the orientation of the handwriting on entomological labels it can not be determined with certainty if the person who prepared the material is right or left-handed. It's well known that many people are strictly one handed in everyday life but show many instances of using the opposite extremity.

> **Julio Antonio Genaro** 25 no. 466, Vedado 12300 Cuidad de La Habana, Cuba



# MORE ON THE SHOCKING ANSWERS TO THE PROBLEM OF STINGS AND BITES

Electric shock therapy (treatment ?) for stings and bites of venomous animals. especially insects, spiders, scorpions and snakes, is becoming more widely appreciated and used (see SPHECOS 25:20). The method of action seemed puzzling to the physicians consulted, however. A Dec. 1986 article in Lancet provides the best approach to understanding the mechanism of action of this novel treatment: Venoms are complex, fragile mixtures of enzymes and their cofactors, along with many other substances. Electrical currents passing through such mixtures could alter the structure and properties of components of the mixture, rendering them incapable of producing their usual deleterious effects. These alterations could result from the destruction of hydrogen bonds between molecules or atoms of enzymes; reduction of some of the metal ions that normally function as cofactors of the enzymes within the venoms; and/ or the current could affect the concerned cellular membranes in the victim, decreasing permeability or transport of the venom within the body. When one considers the difficulties in getting material to study "scientifically" (a bite victim willing to wait for study of individual effects on separate systems before treatment), plus the ethical questions involved in having controls, the tendency to "accept because it works and keep going" is verv understandable.

Copies of the Dec. 1986 Lancet article can be obtained from Terry Nuhn, c/o Sphecos.

Margaret S. Collins Dept. of Entomology Museum of Natural History Smithsonian Institution Washington, D.C. 20560

# EARLIEST PUBLISHED USE OF PHOTOGRAPHS OF HYMENOPTERA?

I recently bought several papers by G. Ernst Adolph on the wings of insects. The earliest one, Ueber Insectenflügel, was published in 1879 in Nova Acta Ksl. Leop.-Carol.-Deutschen Akad. Naturforscher, vol. 41. Four of the six plates

that accompany the work are photomicrographs of wasp and bee wings: Tiphia, Hylaeus, Chrysis, Elampus, Apis mellifica, Nomada, and Sirex. The images are circular and apparently were taken by a camera mounted on a microscope. The wings themselves may have been slide mounted. Does anyone know of an earlier use of actual photographs in a publication on Hymenoptera? More wing photographs appear in an 1883 paper by Adolph titled Zur Morphologie der Hymenopterenflügel. Augleich ein Beitrag zu den Fragen der Species bildung und des Atavismus published in the same journal. These are mostly of honey bee wings.

#### Arnoid S. Menke

# PUZZELMENTIA

Response to a PUZZLEMENTIA from an earlier issue of Sphecos ("INQUIER-ENDA: Why do many stinging wasps have red abdomens?" – 22:18): probably not for the same reason that they have black abdomens (or metasomata or gasters). Actually, the question should have been more or less limited to the Sphecidae, inasmuch as red metasomata are far less common among the Vespidae, Pompilidae, Scoliidae, etc. Even within the Sphecidae it seems to be a <u>common</u> feature largely within the Larrinae, less conspicuous to rare in the remaining subfamilies.

**Roy Snelling** 

### NEW ZEALAND PLACE-NAMES REVISITED

Dear Arnold:

You will be delighted to know that I have just described a lovely, indeed magnificent, new Ammophila with an incredibly long, flutelike "abdomen" (I do so, and I know you will forgive me rather than have its description wait on your long anticipated monograph). It absolutely requires a hippopotomonitrosesquipedalian specific name. And so it now has. I have blessed it as Ammophila philotetaumatawhakatangihangakoauauotamateaturipukakapikimaungahoronukupokaiwhenuakitanata-huensis Mihi. Quite unique, nesting among Sphenodon on North Island, NZ. The name, I am told via an old Maori, is used both geographically, and colloquially when specifying that wasp in subtle allusion to its "abdomen"; I put abdomen in quotes for I am sure he means only the truly spectacular petiole.

But a question for you before submission of the MS: is that name preoccupied?

All the best to you,

# Ken Cooper University of California,

Riverside

PS: It seems to belong in your *procera* group.

# NANCY AND THE CICADA KILLER

Last July your editor was in his backyard talking to his cat, Spook\*, who was sitting on the grass. I noted something moving in the grass behind Spook, and was rather surprised that he had not heard the commotion. Upon closer examination, the moving object turned out to be *Sphecius speciosus* (Drury), the common Cicada Killer. It had a cicada and was struggling to walk with it through the grass. My wife, Nancy, was on the

telephone in the house, and I yelled to her to come out and see something interesting. The phone cord is a long one and she came outside, still talking to her real estate agent. I pointed to the Cicada Killer, and she mentioned to her agent that this huge wasp was in the grass with a cicada. Apparently the wasp was looking for a vertical substrate to gain altitude in order to fly away with her prey. Well, as it turned out, Nancy was a suitable vertical substrate because the Cicada Killer headed right for her left lea! It climbed on to her shoe. Seeing the path it was taking, Nancy grabbed her pant leg tightly, preventing the wasp from taking the inside passage (or scenic route). The wasp proceeded up the outside of her left pant leg. Nancy proceded to describe this activity to her real estate agent who was amazed at Nancy's lack of fear. "You mean you are letting that creature crawl up your body !?" The Cicada Killer continued at a fairly fast pace upward past Nancy's midsection. It was obvious that the wasp would continue until it reached her neck and hair, so Nancy put her left arm across her chest, the Cicada Killer climbed on, and Nancy calmly and slowly outstretched her arm. The wasp walked out to her hand and

took flight with cicada in tow, disappearing over our neighbor's hedge row. We both had a good laugh over this interesting experience. Our Spook, however, seemed completely oblivious to the whole episode.

Spook is so named because I obtained him from the animal shelter on Halloween. He is all black and when I saw him in his cage, all I could see were his two gold eyes peering out at me - that was it, he was a Spook. His name is also a fitting because he is easily spooked by sudden movements or sounds. Well, Spook is no ordinary cat. He weighs 20 pounds and he is so huge that we call him our pet leopard. Some people claim that he is attack-trained.

# BIG BLUE BOOK ERRATA Part 21

- p. 69, LC, L 9: *dahlbomii* is correct, not *dahlbomi*.
- p. 77, LC, L 32: aeneola is correct, not aereola.
- p. 164, LC, L 17 from bottom: precede entry with a dagger symbol (†) and add following to end of entry: nec *Psen reticulata* Cameron, 1902.
- p. 184, RC, L 7: *dahlbomi* Kohl, 1805 is correct entry.
- p. 347, LC, L 27 from bottom: correct distribution is Kenya.



# **RECENT LITERATURE**

(Worth a look: Billen, 1992, Bitsch & Leclercq, 1993, Brothers & Carpenter, 1993, Goulet & Huber, 1992. LaSalle & Gauld, 1993, Sorg, 1988.)

Archer, M.E.

- 1993. Notable wasps and bees (Hym., Aculeata) taken on Jersey and Sark. Ent. Mon. Mag. 128:45-47.
- Alexander, Byron A., Robert L. Minckley and Douglas Yanega
- 1993. Nesting biology of Glenostictia pictifrons (F. Smith) (Hymenoptera: Sphecidae: Bembicini). J. Kansas Ent. Soc. 66(1):108-120.

Antropov, A.V.

- 1993. Alinia carinata gen. et sp. n. a new representative of the tribe Crabronini (Hymenoptera, Sphecidae) from South America. Ent. Obozr. 73 (1):190-193. (in Russian with English summary)
- 1993. A new species of Belomicrus A. Costa (Hymenoptera, Sphecidae) from the Far East. Byullet. Moscov. obshch. ispyt. prirody Otd. Biol. 98 (2):27-28. (in Russian with English summary)

Azevedo, Celso Oliveira

1993. On Rhabdepyris Kieffer (Hymenoptera, Bethylidae) in the area of São Carlos, State of São Paulo, Brazil: II. Subgenus Trichotepyris. Revta bras. Ent. 37(2):305-308.

Archer, Michael E.

- 1993. Notable wasps and bees (Hym., Aculeata) taken on Jersey and Sark. Ent. Mon. Mag. 129:45-47.
- 1993. The life history and colonial characteristics of the homet, Vespa crabro L. (Hym., Vespinae). Ent. Mon. Mag. 129:151-163.
- 1993. The aculeate wasps and bees (Hymenoptera: Aculeata) of Duncombe Park in Watsonian Yorkshire. Naturalist 118:37-44.

Berglind, Sven-Åke

1993. Biotopval och status für vägstekeln Anoplius caviventris (Hymenoptera, Pompilidae) i Sverige. Ent. Tidskr. 114(3):101-1-5.

Billen, Johan

1992. Biology and Evolution of Social Insects. Leuven Univ. Press, Leuven, Belgium, 390 p.

Bitsch, Jacques and Jean Leclercq

1993. Hyménoptères Sphecidae d'Europe Occidentale. Vol. 1. Generalites -Crabroninae. Faune de France 79. 324 p.

Bohart, Richard M.

- 1993, Two new North American species of the tribe Pemphredonini (Hymenoptera: Sphecidae: Pemphredoninae). Pan-Pac. Ent. 69(3):218-220.
- 1993. South American Oxybelus II. The emarginatus group (Hymenoptera, Sphecidae. Insecta Mundi 7(1-2):19-26.
- 1993. South American Oxybelus III. The uniglumis and scutellatus groups (Hymenoptera, Sphecidae. Insecta Mundi 7(1-2):65-76.
- 1993. A synopsis of Central American and Caribbean Oxybelus (Hymenoptera, Sphecidae. Insecta Mundi 7(3):159-168.
- Brothers, Denis J. and James M. Carpenter
- 1993. Phylogeny of Aculeata: Chrysidoidea and Vespoidea. J. Hym. Res. 2 (1):227-302
- Budrys, E.R. and Vladimir L. Kazenas
- 1993. New species of digger wasps of the genus Diodontus (Hymenoptera, Specidae). Ent. Rev. 72(1):114-121.

Carpenter, James M.

1993. Biogeographic patterns in the Vespidae (Hymenoptera): Two views of Africa and South America, Ch. 7, p. 139-155 in: Biological relationships between Africa and South America, Peter Goldbratt (ed.). Yale Univ. Press, New Haven and London.

Clemons, Laurence

1993. Polistes dominulus (Christ, 1791) (Hymenoptera: Vespidae) in greater London. Ent. Record & J. Variation 105:177.

Cölin, Klaus

1993. Bausteine zur Entomofauna des Dorfes - Untersuchungen an ausgewählten Hymenoptera Aculeata und an Syrphidae. Verh. Westd. Entom. Tag 1991:83-90.

Colin, Klaus and Dorothee Behr

1993. Hornissen im Regierungsbezirk Trier. Zusammenstellung von Nachmeldungen und Beobachtungen aus dem Jahr 1992. Dendrocopos 20:142-147. Colomo de Correa, Maria V.

- 1991. Contribucion al conocimiento de los Pompilinae Argentinos III. (Hymenoptera: Pompilidae). Poecilopompilus victori sp.n. Acta Zool. Lilloana 40:53-57.
- 1992. Contribución al conocimiento de los Pompilinae Argentinos (Hymenoptera: Pompilidae). IV. Observaciones sobre algunas especies de Poecilopompilus Howard. Rev. Soc. Ent. Argentina 51(1-4):53-61. Cooper, Martin
- 1993. Two new species of Polybia (Hym., Vespidae) with notes on other members of the genus. Ent. Mon. Mag. 129:31-38.

1993. A new species of Chartergellus (Hym., Vespidae, Polistinae, Polybiini) from Bolivia. Ent. Mon. Mag. 129:165-166. Dani, F.R., R. Cervo and S. Turillazzi

1992. Preliminary observations on abdominal stroking behaviour in Polistes dominulus (Christ) (Hymenoptera: Vespidae), p. 281-285 in: Biology and evolution of social insects, J. Billen, (ed.). Leuven Univ. Press, Leuven, Belgium, 390 p.

Else, G.R.

1993. Recent records of Philanthus triangulum (F.) (Hym., Sphecidae) from southern England. Ent. Mon. Mag. 129:163-164.

Evans, Howard E.

1993. Observations of aggregations of males of two species of beewolves (Hymenoptera: Sphecidae: Philanthus). Psyche 100:25-33.

Field, Jeremy

- 1992. Intraspecific parasitism as an alternative reproductive tactic in nestbuilding wasps and bees. Biol. Rev. 67:79-126.
- 1992. Behavioural differences between English and Japanese populations of Ammophila sabulosa (L.) (Hym., Sphecidae). Ent. Mon. Mag. 128:193-195.
- 1992. Intraspecific parasitism and nest defence in the solitary pompilid wasp Anoplius viaticus (Hymenoptera: Pompilidae). J. Zool., Lond. 228:341-350.
- Francescato, E., S. Turillazzi and M.H. Hansell 1993. Glandular apparatus associated with the gastral tergal tegument in males of Stenogaster concinna and Anischnogaster laticeps (Hymenoptera: Stenogastrinae). Ins. Soc. 40:313-317.

Garcia, Marcos V.B. and Joachim Adis

1993. On the biology of Penepodium goryanum (Lepeletier) in wooden trapnests (Hymenoptera, Sphecidae). Proc. Ent. Soc. Wash. 95(4):547-553. Gaston, K.J.

1993. Spatial patterns in the description and richness of the Hymenoptera, Ch. 12, p. 277-293 in: Hymenoptera and biodiversity, J. LaSalle and I.D. Gauld (eds.). CAB International, Wallingford, Oxon, UK. 348 p.

Genaro, Julio Antonio and Coralia S. Sánchez

- 1992. Observations on the nesting behavior of Tachysphex antillarum (Hymenoptera: Sphecidae). Caribbean J. Sci. 28:218-220.
- 1992. Conducta de nidificacion de Astata unicolor en Cuba (Hymenoptera: Sphecidae). Caribbean J. Sci. 28:220-222.
- 1993. Conducta de nidificacion de Cerceris cerveze, C. cubensis y C. festiva en Cuba (Hymenoptera: Sphecidae). Caribbean J. Sci. 29(1-2):39-43.

Genise, Jorge F. and Lynn S. Kimsey 1993. Revision of the South American thynnine genus Elaphroptera Guérin-Méneville (Hymenoptera: Tiphiidae). J. Hym. Res. 2(1):195-220.

Gess, F.W. and S.K. Gess

1993. Effects of increasing land utilization on species representation and diversity of aculeate wasps and bees in the semi-arid areas of southern Africa, Ch. 4, p. 83-113 in: Hymenoptera and biodiversity, J. LaSalle and I.D. Gauld (eds.). CAB International, Wallingford, Oxon, UK. 348 p.

Giannotti, Edilberto and Cynthia B. Mansur

1993. Dispersion and foundation of new colonies in Polistes versicolor (Hymenoptera, Vespidae). An. Soc. Ent. Brasil 22(2):307-316.

Giannotti, Edilberto and Cláudia Vieira da Silva 1993. Mischocyttarus cassununga (Hymenoptera, Vespidae): external morphology of the brood during the post-embryonic development. Revta bras.

Ent. 37(2):309-312. Gordh, Gordon, Wu Quang Con and E.S. Sugonyaev

1993. Goniozus hanciensis Gordh, sp.n. (Hymenoptera, Bethylidae) - a parasite of the rice leaf roller Gnaphalocrosis medinalis Guenee in North Vietnam, Ent. Obozr. 72:177-185.

Goulet, Henri and John T. Huber, editors

1993. Hymenoptera of the World. An identification guide to families. Centre for Land and Biological Resources Research, Research Branch, Agriculture Canada, Publication 1894/E. Ottawa. 668 p.

Hamon, Jacques

- 1992. Une nouvelle sous-espece de Scoliidae d'Europe occidentale, Scolia (Scolia) hortorum nouveli n.subsp. Sciences Nat, Bull. no. 74:16.
- 1992. Une espèce méconnue de Scoliidae européenne: Colpa klugii (Vander Linden, 1827) (Hymenoptera). L'Entomologiste 48(6):313-316.
- 1992. Observations sur Scolia (Scolia) galbula (Pallas, 1771), Scolia (Scolia) fallax Eversmann, 1849 et Scolia (Discolia) hirta (Schrank, 1781) (Hymenoptera, Scoliidae). Nouv. Revue Ent. (N.S.) 10(1):87-96. Hamon, Jacques and Severiano F. Gayubo

1992. Présence en France et en Espagne de formes mélaniques de Scolia (Scolia) sexmaculata (O.F. Muller, 1766) et S.(S.) hontorum Fabricius, 1787 (Hymenoptera, Scoliidae). Bull. Soc. linn. Bordeaux 20(2):115-123.

Henderson, Gregg, David C. Blouin and Robert L. Jeanne 1993. Yellowjacket (Hymenoptera: Vespidae) repellency by natural products of paper wasps and Avon's Skin-so-Soft<sup>e</sup>, J. Ent. Sci. 28(4):387-392.

Hermann, Henry R., Murray S. Blum and Henry M. Fales 1993. Venom spraying by a tropical polistine wasp (Hymenoptera: Vespidae:

Polistinae). Sociobiology 23:95-99. Hughes, Colin R., David C. Queller, Joan E. Strassmann and Scott K. Davis

1993. Relatedness and altruism in Polistes wasps. Behav. Ecol. 4:128-137.

Jakubzik, Andrea and Klaus Cölin

- 1993. Zur Ökofaunistik Kunstnester bewohnender aculeater Hymenopteren. Verh. Westd. Entorn. Tag 1991:91-100.
- 1993. Zu den Brombeerstengel bewohnenden Hymenopteren (Hymenoptera, Aculeata) von Wehlen (Kreis Bernkastel-Wittlich). Dendrocopos 20:133-141. Jarzembowski, E.A.
- 1991. New insects from the Weald Clay of the Weald. Proc. Geol. Ass. 102 (2):93-108
- Joseph, K.J. and P. Kurian Raphael
- 1991. The nesting and provisioning behaviour of a potter wasp (Eumenidae): an ethological analysis. Entomon 16(1):99-102.
- Karsai, István and Zsolt Pénzes
- 1993. Comb building in social wasps: self-organization and stigmergic script. J. Theor. Biol. 161:505-525.
- Kazenas, Vladimir L.
- 1993. New species of sphecid wasps of the genus Belomicrus A. Costa (Hymenoptera, Sphecidae) from Kazakhstan. Ent. Obozr. 72(1):186-189 (in Russian with English summary)
- 1993. New species of digger wasps of the genus Miscophus (Hymenoptera, Sphecidae) from Kazakhstan. Ent. Rev. 72(3):100-110. (English translation from Zool. Zh. 71(10):97-108, 1992.)
- 1993. New species of Harpactus digger wasps (Hymenoptera, Sphecidae) from Kazakhstan and Uzbekistan. Ent. Rev. 72(6):77-82. (English translation from Ent. Obozr. 72(3):640-645, 1992.)
- Kazenas, Vladimir L. and Byron P. Alexander
- 1993. The nest, prey and larva of Entomosericus kaufmani Radoszkowski (Hymenoptera: Sphecidae). J. Hym. Res. 2(1):221-226.
- Keegans, S.J., E.D. Morgan, B.D. Jackson, G. Berresford, S. Turillazzi and J. Billen
- 1992. The chemical nature and origin of the abdominal substance of the social wasp Liostenogaster flavolineata (Vespidae, Stenogastrinae), p. 217-221 in: Biology and evolution of social insects, J. Billen, (ed.). Leuven Univ. Press, Leuven, Belgium. 390 p.
- Kilimnik, A.N.
- 1993. Chrysidid wasps of the genus Hedychrum Latreille, 1806 (Hymenoptera, Chrysididae) of East Europe. Ent. Obozr. 72(2): 388-400. (in Russian with English summary)
- Kimsey, Lynn S.
- 1993. New neotropical Amisegine wasps (Hymenoptera: Chrysididae). Pan-Pac. Ent. 69(3):205-212.
- 1993. An unusual new Tiphild genus from Peru and a key to the American genera of Tiphiinae (Hymenoptera). Pan-Pac. Ent. 69(3):213-217.
- 1993. Review of the sphecid genus Paradolichurus, and description of a new species (Hymenoptera, Sphecidae). J. Kansas Ent. Soc. 66(2):245-249. Kojima, Jun-ichi
- 1993. Feeding of larvae by males of an Australian paper wasp, Ropalidia plebeiana Richards (Hymenoptera, Vespidae). Jpn. J. Ent., 61(2):213-215. Kurian Raphael, P. and K.J. Joseph
- 1993. Bionomics of the mason wasp Xenorhynchium nitidulum (Fabr.) (Eumenidae) with special reference to mating behaviour. Proc. Kerala Sci. Congress, Feb.-Mar. 1991, Kozhikode, pp. 360-361.

LaSalle, J. and I.D. Gauld

- 1993. Hymenoptera: their diversity and their impact on the diversity of other organisms, Ch. 1, p. 1-26 in: Hymenoptera and biodiversity, J. LaSalle and I.D. Gauld (eds.). CAB International, Wallingford, Oxon, UK. 348 p.
- Leclercq, Jean
  - 1993. Une espece nouvelle d'Odontocrabro, du Maroc. Lambillionea 93(2): 191-192.
- Leclercq, Jean
- 1993. Atlas de répartition des Crabroniens de France et des régions limitrophes (Hymenoptera, Sphecidae, Crabronini). Notes Fauniques de Gembloux (27):3-94.
- Lelej, Arkady S.
- 1993. Phylogeny and evolution of Mutillidae and their position in Scolioidea (Hymenoptera). Doctoral dissertation, Zool. Inst. Russian Akad. Sci. 42 p. Li Qiang and Yang Chi-kun
- 1992. Three new species of Podalonia from China (Hymenoptera: Sphecidae). Acta Agric. Pekinensis 18:85-90. (in Chinese with English summary) Lorenzi, M.C. and R. Cervo
- 1992. Behaviour of Polistes biglumis bimaculatus (Hymenoptera, Vespidae) foundresses on alien conspecific nests, p. 273-279 in: Biology and evolution of social insects, J. Billen, (ed.). Leuven Univ. Press, Leuven, Belaium, 390 p.
- Luft. Patrick A.
- 1993. Experience affects oviposition in Goniozus nigrifernur (Hymenoptera: Bethylidae). Ann. Ent. Soc. Amer. 86(4):497-505.
- Melo, Gabriel Augusto R. de and Lucio Antonio de O. Campos
- 1993. Nesting biology of Microstigmus myersi Turner, a wasp with longhaired larvae (Hymenoptera: Sphecidae, Pemphredoninae). J. Hym. Res. 2(1):183-188.

Moya-Raygoza, Gustavo and Javier Trujillo-Arriaga

- 1993. Evolutionary relationships between Dalbulus leafhopper (Homoptera: Cicadellidae) and its dryinid (Hymenoptera: Dryinidae) parasitoids. J. Kansas Ent. Soc. 66(1):41-50.
- Noguchi, Shizuo
- 1993. Provisioning behaviour of Delta esuriens. Gekkan-Mushi. 271:40-41. (in Japanese)
- Ohl, Michael
- 1993. Morphologische Untersuchungen am Exoskelett der Grabwespen als Grundlage für die Ermittlung der phylogenetischen Beziehungen innerhalb der Sphecidae (Hymenoptera). Masters thesis, Mathematische-Naturwissenschlaftlich Fakultat der Christian-Albrechts-Universitat, Kiel. 167 p.
- Pénzes, Zsolt and István Karsai
- 1993, Round shape produced by stigmergic scripts in social wasp. Proc. European Conf. Artificial Life (Brussels), p. 896-905.
- Petit, J.
- 1993. Faunistique de la Montagne Saint-Pierre. 2. Présence de Discoelius zonata (Panzer) (Hymenoptera Eumenidae). Lambillionea 93(2):139-142. Piek, Tom
- 1993. New neurotoxins from venom of aculeate Hymenoptera: a contribution to the knowledge of stinging behavior. J. Hym. Res. 2(1):101-105.
- Queller, David C.
- 1992. Quantitative genetics, inclusive fitness, and group selection. Amer. Nat. 139(3):540-558
- 1992. A general model for kin selection. Evolution 46(2):376-380.
- 1992. Does population viscosity promote kin selection? Trends Ecol. Evol. 7 (10):322-324.
- 1993. Worker control of sex ratios and selection for extreme multiple mating by queens. Amer. Nat. 142(2):346-351.
- Queller, David C., Joan E. Strassmann and Colin R. Hughes
- 1993. Microsatellites and kinship. Trends Ecol. Evol. 8(8):285-288.
- 1993. Microsatellite markers. Trends Ecol. Evol. 8(8):?
- Queller, David C., Jorge A. Negrón-Sotomayor, Joan E. Strassmann and Colin R. Hughes
- 1993. Queen number and genetic relatedness in a neotropical wasp, Polybia occidentalis. Behav. Ecol. 4:7-13.
- Queller, David C., Joan E. Strassmann, Carlos R. Solís, Colin R. Hughes and Debbie Moralez DeLoach.
- 1993. A selfish strategy of social insect workers that promotes social cohesion. Nature 365:639-641.
- Strassmann, Joan E.
- 1993. Weak gueen of social contract? Nature 363:502-503.
- Rasnitsyn, Alexandr P.
- 1993, Archaeoscolinae, an extinct subfamily of scoliid wasps (insecta: Vesppida = Hymenoptera: Scoliidae). J. Hym. Res. 2(1):85-96. Schwert, Donald P.
- 1993. Book Review: Life in Amber, by G.O. Poiner, Jr. Ann. Ent. Soc. Amer. 86(4):509-510.
- Silvera, Orlando Tobias and Flavio Henrique Caetano
- 1993. A morphometric study of sting glands in vespid wasps (Hymenoptera, Vespidae). Sociobiology 23(1):45-62.
- Soika, Antonio Giordani
  - 1993. Controbuto alla conoscenza delgli Eumenidi dell'Australia e della Nuova Guinea. Boll. Mus. civ. St. nat. Venezia 42(1991):125-149.
  - 1993. Di alcuni nuovi Eumenidi della Regione Orientale (Hym. Vespidae). Boll. Mus. civ. St. nat. Venezia 42(1991):151-163.
- Sorg, Martin
- 1988. Zur phylogenie und systematik der Bethylidae (Insecta: Hymenoptera: Chrysidoidea). Geol. Inst. Univ. Koln Sonderveroeffentlich. 63:1-146.

Sorg, Martin and Klaus Colin

- 1993. Die Wegwespen (Hymenoptera, Pompilidae) von Gönnersdorf (Kr. Daun). Beigräge zur Insektenfaune der Eifeldörfer XI. Dendrocopos 20:161-169. Spradbery, J.P.
  - 1993. Queen brood reared in worker cells by the social wasp, Vespula germanica (F.) (Hymenoptera: Vespidae). Ins. Soc. 40:181-190.
- Suzuki, T.
- 1993. Timing of insemination in terms of age of new reproductive females of a temperate-zone polistine wasp, Polistes snelleni (Hymenoptera: Vespidae). ins. Soc. 40:341-343.
- Turillazzi, S.

1992. Nest usurpation and social parasitism in Polistes wasps: new acquisitions and current problems, p. 263-272 in: Biology and evolution of social insects, J. Billen, (ed.). Leuven Univ. Press, Leuven, Belgium. 390 p. Willink, Abraham and Arturo Roig Alsina

1993. On Odynerus rachiphorus Schletterer, a Masarinae (Trimeria), not a Eumeninae (Hymenoptera, Vespidae). J. Hym. Res. 2(1):303.

Yu. Dicky S. 1993. A proposed system for stabilizing names of species, illustrated with reference to the Ichneumonidae (Hymenoptera). Bull. Zool. Nomen. 50(1):7-11.